

# Publications mathématiques de Besançon

## ALGÈBRE ET THÉORIE DES NOMBRES

Yuichiro Hoshi

### Introduction to Mono-anabelian Geometry

2021, p. 5-44.

<https://doi.org/10.5802/pmb.42>

© Les auteurs, 2021.



Cet article est mis à disposition selon les termes de la licence  
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL.  
<https://creativecommons.org/licenses/by/4.0/deed.fr>

L'accès aux articles de la revue « Publications mathématiques de Besançon »  
(<http://pmb.centre-mersenne.org/>) implique l'accord avec les conditions générales  
d'utilisation (<http://pmb.centre-mersenne.org/legal/>).

*Publication éditée par le laboratoire de mathématiques  
de Besançon, UMR 6623 CNRS/UFC*



*Les Publications mathématiques de Besançon sont membres du  
Centre Mersenne pour l'édition scientifique ouverte  
<http://www.centre-mersenne.org/>*

# INTRODUCTION TO MONO-ABELIAN GEOMETRY

*by*

Yuichiro Hoshi

---

*Abstract.* — The present article is based on the four hours mini-courses “Introduction to Mono-anabelian Geometry” which the author gave at the conference “Fundamental Groups in Arithmetic Geometry” (Paris, 2016). The purpose of the present article is to introduce *mono-anabelian geometry* by focusing on mono-anabelian geometry for *mixed-characteristic local fields*, which provides elementary but nontrivial examples of typical arguments in the study of mono-anabelian geometry.

*Résumé.* — (*Introduction à la géométrie mono-anabélienne*) Cet article est basé sur les 4 heures de mini-cours « Introduction to Mono-anabelian Geometry » que l’auteur a données lors de la conférence « Fundamental Groups in Arithmetic Geometry » (Paris, 2016). L’objectif est de présenter la géométrie mono-anabélienne en se concentrant sur les corps locaux de caractéristique mixte ce qui permet de fournir des exemples élémentaires mais non-triviaux du type d’arguments présents dans l’étude de géométrie mono-anabélienne.

## Introduction

The present article is based on the four hours mini-courses “Introduction to Mono-anabelian Geometry” which the author gave at the conference “Fundamental Groups in Arithmetic Geometry” (Paris, 2016). In the present article, we discuss *mono-anabelian geometry*.

*Anabelian geometry* is, in a word, an area of arithmetic geometry in which one studies the geometry of geometric objects of interest from the point of view of arithmetic fundamental groups. Put another way, roughly speaking, anabelian geometry discusses the issue of how much information concerning the geometry of geometric objects of interest is contained in the knowledge of the arithmetic fundamental groups.

The classical point of view of anabelian geometry (i.e., more precisely, of *Grothendieck’s anabelian conjecture*) centers around a *comparison* between *two* geometric objects of interest

---

**2020 Mathematics Subject Classification.** — 11S20.

*Key words and phrases.* — mono-anabelian geometry, MLF, mono-anabelian reconstruction algorithm, MLF-pair, cyclotomic synchronization, Kummer poly-isomorphism, mono-anabelian transport.

*Acknowledgements.* — The author was supported by the Inamori Foundation and JSPS KAKENHI Grant Number 15K04780.

via the arithmetic fundamental groups. In fact, in a discussion of anabelian geometry, typically, one fixes *two* geometric objects and discusses the relationship between a certain set of morphisms (e.g., the set of isomorphisms) between the fixed two objects and a certain set of homomorphisms (e.g., the set of isomorphisms) between the étale fundamental groups. In particular, roughly speaking, the classical point of view of anabelian geometry may be summarized as the study of some properties such as *faithfulness/fullness* of the (restriction, to a certain suitable category of geometric objects, of the) functor of taking arithmetic fundamental groups. Moreover, the term "*group-theoretic*" (that often appears in discussions of anabelian geometry) is, in the classical point of view, defined simply to mean "*preserved by an arbitrary isomorphism between the arithmetic fundamental groups under consideration*". In [6], this classical point of view is referred to as "*bi-anabelian geometry*".

bi-anabelian geometry

$$\pi_1^{\text{ét}}(X) - \pi_1^{\text{ét}}(X \cdot) \stackrel{?}{=} \text{(objects related to)} X - \text{(objects related to)} X.$$

By contrast, *mono-anabelian geometry* centers around the task of establishing a "*group-theoretic software*" (i.e., "*group-theoretic algorithm*") whose input data consists of a *single abstract topological group* isomorphic to the arithmetic fundamental group of a *single* geometric object of interest.

In particular, a *mono-anabelian reconstruction algorithm* (i.e., a "group-theoretic algorithm" discussed in mono-anabelian geometry) has the virtue of being *free* of any mention of some "fixed reference model" copy of geometric objects (as the above " $X$ " for " $\pi_1^{\text{ét}}(X)$ " in the case of bi-anabelian geometry). In the point of view of mono-anabelian geometry, the term "*group-theoretic algorithm*" is used to mean that "*the algorithm in a discussion is phrased in language that only depends on the topological group structure of the arithmetic fundamental group under consideration*" (cf., e.g., [6, Introduction, Remark 1.9.8, and Remarks following Corollary 3.7] for more details concerning bi-anabelian/mono-anabelian geometry).

mono-anabelian geometry

$$\text{a topological group isomorphic to } \pi_1^{\text{ét}}(X)$$

?

object(s) isomorphic to (objects related to)  $X$

The purpose of the present article is to introduce *mono-anabelian geometry* by focusing on mono-anabelian geometry for *mixed-characteristic local fields*, i.e., *MLF's* (cf. Definition 1.1), which provides elementary but nontrivial examples of typical arguments in the study of mono-anabelian geometry.

In Section 1, we introduce some notational conventions related to MLF's and recall some basic facts concerning objects that arise from MLF's. These basic facts will be applied in other sections of the present article. In Section 2, we recall some results in *bi-anabelian geometry* for MLF's. Some of the fundamental results in bi-anabelian geometry for MLF's are as follows (cf. Theorem 2.2):

- The isomorphism class of an MLF is *not determined* by the isomorphism class of the absolute Galois group of the MLF.
- There exists an outer automorphism of the absolute Galois group of an MLF which does *not arise* from any automorphism of the MLF.

These results lead us to an interest in a study of *conditions* for an outer isomorphism between the absolute Galois groups of MLF's to arise from an isomorphism between the original MLF's. In Section 2, we also recall such *conditions* (cf. Theorem 2.3, Remark 2.3.2).

In Sections 3 and 4, we establish some *mono-anabelian reconstruction algorithms* for MLF's, i.e., some "group-theoretic algorithms" whose input data consist of a group of *MLF-type* (i.e., an abstract group *isomorphic to the absolute Galois group of an MLF* — cf. Definition 3.1). For instance, by applying the mono-anabelian reconstruction algorithms discussed in Section 4, one may construct, from a group  $G$  of MLF-type,  $G$ -monoids

$$\bar{O}^\times(G) \quad \bar{O}^B(G) \quad \bar{k}^\times(G)$$

which "correspond" to the  $\text{Gal}(\bar{k}/k)$ -monoids

$$O_k^\times \quad O_k^B \quad \bar{k}^\times,$$

where  $k$  is an MLF, and  $\bar{k}$  is an algebraic closure of  $k$  (i.e.,

- the multiplicative  $\text{Gal}(\bar{k}/k)$ -module  $O_k^\times$  of units of the ring of integers of  $\bar{k}$ ,
- the multiplicative  $\text{Gal}(\bar{k}/k)$ -monoid  $O_k^B$  of nonzero integers of  $\bar{k}$ , and
- the multiplicative  $\text{Gal}(\bar{k}/k)$ -module  $\bar{k}^\times$  of nonzero elements of  $k$ ,

respectively (cf. Summary 3.15, Summary 4.3).

One important aspect of mono-anabelian geometry is the technique of *mono-anabelian transport*. In order to explain mono-anabelian transport, in Section 5, we introduce the notion of an *MLF-pair* (cf. Definition 5.3). Some types of MLF-pairs are discussed in the present article. For instance, an *MLF<sup>×</sup>-pair* is defined to be a collection of data  $G \curvearrowright M$  consisting of a monoid  $M$ , a group  $G$ , and an action of  $G$  on  $M$  such that there exists an isomorphism (in the evident sense) of  $G \curvearrowright M$  with the collection of data

$$\text{Gal}(\bar{k}/k) \curvearrowright O_k^\times$$

for some MLF  $k$  and some algebraic closure  $\bar{k}$  of  $k$ . If  $G \curvearrowright M$  is an MLF-pair, then we shall refer to the group  $G$ , the monoid  $M$  as the *étale-like portion*, the *Frobenius-like portion* of  $G \curvearrowright M$ , respectively (cf. Definition 5.4).

In Section 6, we discuss a phenomenon of *cyclotomic synchronization* for MLF-pairs. Let us recall that a *cyclotome* refers to an "object isomorphic to the object  $Z(1)$ ". Various (a priori independent) cyclotomes often appear in studies of arithmetic geometry. For instance, if  $\bar{k}$  is an algebraically closed field of characteristic zero, then each of

- the cyclotome

$$(\bar{k}) \stackrel{\text{def}}{=} \varprojlim_n \text{Ker}(\bar{k}^\times \xrightarrow{\times n} \bar{k}^\times)$$

— where the projective limit is taken over the positive integers  $n$  — associated to  $\bar{k}$  and

- the dual

$$(C) \stackrel{\text{def}}{=} \text{Hom}_Z H_{\text{ét}}^2(C, Z), Z$$

over  $Z$  of the second étale cohomology  $H_{\text{ét}}^2(C, Z)$  of a projective smooth curve  $C$  over

gives an example of a cyclotome.

If one works with certain scheme/ring structures of objects related to cyclotomes under consideration, then one may obtain a phenomenon of *cyclotomic synchronization*, i.e., synchronization of cyclotomes. For instance, in the case of the above examples ( ) and (C), the homomorphism  $\text{Pic}(C) \rightarrow H^2(C, \mathbb{Z})$  obtained by considering the *first Chern classes* yields an isomorphism

$$\text{Pic}(C)/\text{Pic}^0(C) \xrightarrow{\sim} \mathbb{Z} \cong \text{Hom}_{\mathbb{Z}}(\mathbb{Z}, \mathbb{Z});$$

thus, an invertible sheaf on  $C$  of *degree one* determines a *cyclotomic synchronization* (i.e., an isomorphism between cyclotomes) (C)  $\xrightarrow{\sim}$  ( ) by means of which one usually (i.e., in the usual point of view of arithmetic geometry) *identifies* (C) with ( ). On the other hand, observe that such a phenomenon of cyclotomic synchronization usually *depends*, at least in an a priori sense, on a certain scheme/ring structures of objects related to cyclotomes under consideration. Thus, if one works in a situation in which objects related to cyclotomes under consideration *lose* a certain portion of the rigidity that arises from scheme/ring structures, then one *cannot apply* the “scheme/ring-theoretic” construction of the cyclotomic synchronization.

In our study of MLF-pairs, one may construct, from a *single* MLF-pair, (a priori independent) *two cyclotomes*, i.e., a cyclotome constructed from the *étale-like* portion and a cyclotome constructed from the *Frobenius-like* portion (cf. Definition 5.9, Proposition 5.10). In Section 6, we establish a cyclotomic synchronization which relates the Frobenius-like cyclotome to the étale-like cyclotome (cf. Definition 6.6, Proposition 6.7).

In Section 7, we discuss *Kummer poly-isomorphisms* and *mono-anabelian transport*. A *Kummer poly-isomorphism* relates (monoids constructed, via some functorial algorithms, from) *Frobenius-like portions* to *mono-anabelian étale-like monoids* (i.e., monoids constructed, via some mono-anabelian reconstruction algorithms, from *étale-like portions*). In Section 7, we establish Kummer poly-isomorphisms for MLF-pairs (cf. Definition 7.4). Finally, we discuss the technique of mono-anabelian transport (cf. Remark 7.6.1).

Some details of discussions given in the portion from Sections 3 to 7 of the present article may be found in, for instance, [4, §1, §2], [6, §3, §5], [7, §2], [2, §1].

**Acknowledgments.** — The present article is based on the mini-courses which the author gave at the conference “Fundamental Groups in Arithmetic Geometry”. The author would like to thank the organizers of the conference for the time and effort they spent to make the conference a success. The author would like to thank especially *Anna Cadoret*, who is one of the organizers, for inviting me to the conference. The author also would like to thank some participants of the conference for their interest in and some discussions concerning the mini-courses. Finally, the author would like to thank the *referee* for some helpful comments.

## 0. Notational Conventions

*Sets.* — If  $S$  is a finite set, then we shall write  $\#S$  for the *cardinality* of  $S$ . If  $G$  is a group, and  $T$  is a set equipped with an action of  $G$ , then we shall write  $T^G \subset T$  for the subset of  $G$ -invariants of  $T$ .

**Monoids.** In the present article, every monoid is assumed to be commutative. Let  $M$  be a monoid. (The monoid operation of  $M$  will be written multiplicatively). We shall write  $M^\times$  for the abelian group of invertible elements of  $M$ . We shall write  $M^{gp}$  for the groupification of  $M$  (i.e., the abelian group obtained by forming the monoid of equivalence classes with respect to the relation on  $M$  defined by, for  $(a_1; b_1), (a_2; b_2) \in M \times M$ ,  $(a_1; b_1) \sim (a_2; b_2)$  if there exists an element  $c \in M$  of  $M$  such that  $ca_1b_2 = ca_2b_1$ ). We shall write  $M^{pf}$  for the perfection of  $M$  (i.e., the monoid obtained by forming the injective limit of the injective system of monoids

$$! \ M \ ! \ M \ !$$

given by assigning to each positive integer a copy of  $M$ , which we denote by  $I_n$ , and to each two positive integers  $n, m$  such that  $n$  divides  $m$  the homomorphism  $I_n \rightarrow I_m$  given by multiplication by  $m/n$ ). We shall write  $M^\sim \stackrel{\text{def}}{=} M [f_M g]$ ; we regard  $M^\sim$  as a monoid (that contains  $M$  as a submonoid) by setting  $f_M = g_M \stackrel{\text{def}}{=} M$  and  $a_M \stackrel{\text{def}}{=} M$  for every  $a \in M$ . We shall write  $\mathbb{N}$  for the additive monoid of nonnegative integers.

**Modules** Let  $M$  be a module. If  $n$  is a positive integer, then we shall write  $M[n]$  for the submodule obtained by forming the kernel of the endomorphism of  $M$  given by multiplication by  $n$ . We shall write  $M_{\text{tor}} \stackrel{\text{def}}{=} \bigcup_{n \geq 1} M[n]$  for the submodule of torsion elements of  $M$  and

$$M^\wedge \stackrel{\text{def}}{=} \varprojlim_n M/(nM);$$

where the projective limit is taken over the positive integers  $n$ . (So if  $M$  is finitely generated, then  $M^\wedge$  is nothing but the profinite completion of  $M$ .)

**Topological Groups** Let  $G$  be a topological group. Then we shall write  $G^{ab}$  for the abelianization of  $G$  (i.e., the quotient of  $G$  by the closure of the commutator subgroup of  $G$ ),  $G^{ab\text{-tor}} \stackrel{\text{def}}{=} (G^{ab})_{\text{tor}}$ , and  $G^{ab=\text{tor}}$  for the quotient of  $G^{ab}$  by the closure of  $G^{ab\text{-tor}}$ . Suppose that  $G$  is a profinite group. Then we shall say that  $G$  is slim if, for every open subgroup  $H \leq G$  of  $G$ , the centralizer of  $H$  in  $G$  is trivial. If  $n$  is a nonnegative integer, and  $M$  is a topological  $G$ -module, then we shall write  $H^n(G; M)$  for the  $n$ -th continuous group cohomology of  $G$  with coefficients in  $M$  and

$$\varinjlim_{H \leq G} H^n(G; M) \stackrel{\text{def}}{=} \varinjlim_{H \leq G} H^n(H; M);$$

where the injective limit is taken over the open subgroups  $H \leq G$  of  $G$ .

**Rings.** In the present article, every ring is assumed to be unital, associative, and commutative. If  $R$  is a ring, then we shall write  $R_+$  for the underlying additive module of  $R$  and  $R^\times$  for the multiplicative module of units of  $R$ . If  $R$  is an integral domain, then we shall write  $R^B \leq R$  for the multiplicative monoid of nonzero elements of  $R$ .

**Fields.** Let  $K$  be a field. Then we shall write  $(K)^\times \stackrel{\text{def}}{=} (K)_{\text{tor}}$  for the group of roots of unity in  $K$  and  $K^\times = K \setminus \{0\}$  for the underlying multiplicative monoid of  $K$ . (So we have a natural isomorphism  $(K)^\times \cong K^\times$  of monoids that maps  $\kappa \in (K)^\times$  to  $0 \in K^\times$ ). If,

moreover,  $K$  is algebraically closed and of characteristic zero, then we shall write

$$(K) \stackrel{\text{def}}{=} \varprojlim_n (K)[n] = \varprojlim_n K[n]$$

(where the projective limits are taken over the positive integers) and refer to  $(K)$  as the cyclotome associated to  $K$ . Thus, the cyclotome has a natural structure of profinite, hence also topological, module and is, as an abstract topological module, isomorphic to  $\hat{\mathbb{Z}}_+$ .

**Categories** Let  $A$  be an object of a category. Then we shall refer to an object of the category isomorphic to  $A$  as an isomorph of  $A$ .

Let  $A, B,$  and  $C$  be objects of a category. Then we shall refer to a nonempty set of isomorphisms from  $A$  to  $B$  in the category as a poly-isomorphism (from  $A$  to  $B$ ). (So one may regard a single isomorphism as a poly-isomorphism, i.e., of cardinality one.) Let  $f: A \rightarrow B$  be a poly-isomorphism (i.e., from  $A$  to  $B$ ) and  $g: B \rightarrow C$  a poly-isomorphism (i.e., from  $B$  to  $C$ ). We shall write  $g \circ f: A \rightarrow C$  for the poly-isomorphism (i.e., from  $A$  to  $C$ ) obtained by forming the set  $\{g \circ f_j \mid f_j \in f\}$ ;  $g \circ f$  and refer to  $g \circ f$  as the composite of  $f$  and  $g$ . We shall write  $f^{-1}: B \rightarrow A$  for the poly-isomorphism (i.e., from  $B$  to  $A$ ) obtained by forming the set  $\{f_j^{-1} \mid f_j \in f\}$  and refer to  $f^{-1}$  as the inverse of  $f$ .

## 1. Generalities on MLF's

In the present section, let us introduce some notational conventions related to mixed-characteristic local fields, i.e., MLF's (cf. Definition 1.1 below), and recall some basic facts concerning objects that arise from MLF's. These basic facts will be applied in other sections of the present article.

**Definition 1.1** . We shall refer to a finite extension of  $\mathbb{Q}_p$ , for some prime number  $p$ , as an MLF. Here, MLF is to be understood as an abbreviation for mixed-characteristic local field.

In the remainder of the present section, let  $k$  be an MLF. Then we shall write

$\mathcal{O}_k$  for the ring of integers of  $k$ ,

$\mathfrak{m}_k \subset \mathcal{O}_k$  for the maximal ideal of  $\mathcal{O}_k$ ,

$\underline{k} \stackrel{\text{def}}{=} \mathcal{O}_k / \mathfrak{m}_k$  for the residue field of  $\mathcal{O}_k$ ,

$\mathcal{O}_k^{(n)} \stackrel{\text{def}}{=} 1 + \mathfrak{m}_k^n \subset \mathcal{O}_k$  (where  $n$  is a positive integer) for the  $n$ -th higher unit group of  $k$ ,

$\mathcal{O}_k^{\times} \stackrel{\text{def}}{=} \mathcal{O}_k^{\times} \setminus \mathfrak{m}_k$  for the group of principal units of  $k$ ,

$\mu_k$  for the (uniquely determined) Haar measure on (the locally compact topological module)  $k_+$  such that  $\mu_k((\mathcal{O}_k)_+) = 1$ , and

$\rho_k \stackrel{\text{def}}{=} \text{char}(\underline{k})$  for the residue characteristic of  $k$ .

Thus, one verifies easily that  $[k : \mathbb{Q}_{p_k}]$  and  $[k : \mathbb{F}_{p_k}]$  are finite. We shall write

$$d_k \stackrel{\text{def}}{=} [k : \mathbb{Q}_{p_k}] \text{ and}$$

$$f_k \stackrel{\text{def}}{=} [k : \mathbb{F}_{p_k}].$$

We shall write, moreover,

$$e_k \stackrel{\text{def}}{=} v_k(k^{-1} \mathbb{Z}) \text{ for the absolute ramification index of } k,$$

$$\log_k : \mathbb{Z} \rightarrow \mathbb{Z}_+ \text{ for the } p_k\text{-adic logarithm, and}$$

$$I_k \stackrel{\text{def}}{=} (2p_k)^{-1} \log_k(\mathbb{Z}) \text{ for the log-shell of } k.$$

In the following lemma, let us recall some basic facts concerning the above objects.

Lemma 1.2 . The following hold:

(i) The topological module  $\mathbb{Z}_+$  is isomorphic to the topological module

$$\mathbb{Z}_+ \cong (p_k^{f_k} \mathbb{Z}) \oplus (\mathbb{Z}_+ / p_k^a \mathbb{Z}_+) \oplus (\mathbb{Z}_{p_k})_+^{d_k} \oplus \mathbb{Z}_+$$

for some nonnegative integers. Moreover, the topological submodule  $\mathbb{O}_k$  of  $\mathbb{Z}_+$  corresponds relative to such an isomorphism, to the kernel of the fourth projection

$$\mathbb{Z}_+ \cong (p_k^{f_k} \mathbb{Z}) \oplus (\mathbb{Z}_+ / p_k^a \mathbb{Z}_+) \oplus (\mathbb{Z}_{p_k})_+^{d_k} \oplus \mathbb{Z}_+ \rightarrow \mathbb{Z}_+ :$$

(ii) The topological submodule  $\mathbb{O}_k$  of  $\mathbb{O}_k$  is the maximal pro- $p_k$  submodule of  $\mathbb{O}_k$ .

(iii) It holds that  $d_k = f_k e_k$ .

(iv) The  $p_k$ -adic logarithm  $\log_k : \mathbb{Z} \rightarrow \mathbb{Z}_+$  determines an isomorphism of topological modules

$$(\mathbb{Z})^{p_k} \rightarrow \mathbb{Z}_+ :$$

(v) It holds that  $\text{Ker}(\log_k) = I_k$ .

(vi) It holds that  $(\mathbb{O}_k)_+ \cong I_k$ .

Proof. Assertions (i), (ii) follow immediately from [8, Chapter II, Proposition 5.3 and Proposition 5.7 (i)]. Assertion (iii) follows from [8, Chapter II, Proposition 6.8]. Assertions (iv), (v) follow immediately from [8, Chapter II, Proposition 5.5], together with assertion (i). Assertion (vi) follows immediately from [8, Chapter II, Proposition 5.5], together with the (easily verified) fact that  $e_k > e_k - (p_k - 1)$  (respectively,  $2e_k > e_k - (p_k - 1)$ ) if  $p_k \neq 2$  (respectively,  $p_k = 2$ ).

Next, let us recall some basic facts concerning the measure  $\mu_k$ .



Lemma 1.3 . The following hold:

- (i) Let  $S, T \subseteq k_+$  be compact opensubsets of  $k_+$ . Then the measure  $\mu_k$  satisfies the following conditions:
  - (a) If  $S \cap T = \emptyset$ , then it holds that  $\mu_k(S \cup T) = \mu_k(S) + \mu_k(T)$ .
  - (b) For each  $a \in k_+$ , it holds that  $\mu_k(S + a) = \mu_k(S)$ .
  - (c) If  $S$  is contained in  $O_k \setminus \{0\}$ , and the natural surjection  $S \xrightarrow{\log_k} \log_k(S)$  determined by  $\log_k$  is bijective, then it holds that  $\mu_k(\log_k(S)) = \mu_k(S)$ .

(ii) It holds that

$$\mu_k(O_k) = 1 \quad \mu_k^{f_k} = \mu_k^{f_k} (\mu_k^{f_k})^{-1};$$

(iii) It holds that

$$\mu_k(I_k) = \mu_k^{d_k} (\mu_k^{f_k})^{-1} (\mu_k)^{(p_k)};$$

where we write

$$\mu_k \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } p_k \neq 2 \\ 2 & \text{if } p_k = 2 \end{cases}$$

and  $(\mu_k)^{(p_k)}$  for the  $p_k$ -Sylow subgroup of (the finite cf. Lemma 1.2(i)) abelian group  $(\mu_k)$ .

Proof. First, we verify assertion (i). The assertion that  $\mu_k$  satisfies conditions (a), (b) follows from the definition of a Haar measure. Next, we verify the assertion that  $\mu_k$  satisfies condition (c). Let us recall that the system  $\{O_k^{-n}g_{n-1}\}$  forms a basis of neighborhoods of the identity element  $1 \in O_k$  of  $O_k$  (cf. the discussion preceding [8, Chapter II, Proposition 3.10]). Thus, it follows immediately in light of the (easily verified) fact that the endomorphism of  $k_+$  given by multiplication by an element of  $O_k$  is a topological automorphism which restricts to an automorphism of  $(O_k)_+ \subseteq k_+$  from conditions (a), (b) that, to verify the assertion that  $\mu_k$  satisfies condition (c), it suffices to verify that there exists a positive integer  $n_0$  such that if  $n > n_0$ , then

$$\mu_k(\log_k(O_k^{-n})) = \mu_k(O_k^{-n});$$

On the other hand, it follows from [8, Chapter II, Proposition 5.5], together with condition (b), that if  $n > e_k = (p_k - 1)$ , then

$$\mu_k(\log_k(O_k^{-n})) = \mu_k(m_k^n) = \mu_k(1 + m_k^n) = \mu_k(O_k^{-n});$$

This completes the proof of the assertion that  $\mu_k$  satisfies condition (c), hence also of assertion (i).

Next, we verify assertion (ii). Since (one verifies easily that)  $O_k = (O_k)_+ \cup m_k$ , it follows from condition (a) of assertion (i) that

$$\mu_k(O_k) = \mu_k((O_k)_+) + \mu_k(m_k) = \mu_k((O_k)_+) (1 + [(O_k)_+ : m_k]^{-1}) = 1 + \mu_k^{f_k};$$

as desired. This completes the proof of assertion (ii).

Finally, we verify assertion (iii). Let us first recall from Lemma 1.2(v) that  $\text{Ker}(\log_k) = (\mu_k)$ . Thus, it follows from conditions (a), (c) of assertion (i) that

$$\mu_k(I_k) = [\mu_k : \log_k(O_k)] \mu_k(\log(O_k)) = [\mu_k : \log_k(O_k)] (\mu_k)^{-1} \mu_k(O_k);$$

In particular, since

$$] (k) = ] (k)^{(p_k)} (p_k^{f_k} - 1)$$

(cf. Lemma 1.2(i)), it follows from assertion (ii) and Lemma 1.2(iv) that

$${}_k(I_k) = p_k^{k d_k} ] (k)^{(p_k)}^{-1} (p_k^{f_k} - 1)^{-1} p_k^{f_k} (p_k^{f_k} - 1) = p_k^{k d_k} p_k^{f_k} ] (k)^{(p_k)};$$

as desired. This completes the proof of assertion (iii).

Next, let  $\bar{k}$  be an algebraic closure of  $k$ . Then we shall write

$O_k$   $\bar{k}$  for the ring of integers of  $\bar{k}$ ,

$\bar{k}$  for the residue field of  $O_k$ ,

$\log_k : O_k \rightarrow \bar{k}_+$  for the  $p_k$ -adic logarithm,

$G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=k)$  for the absolute Galois group of  $k$  with respect to  $\bar{k}$ ,

$I_k \subset G_k$  for the inertia subgroup of  $G_k$ , and

$P_k \subset I_k$  for the wild inertia subgroup of  $G_k$ .

Thus,  $\bar{k}$  is an algebraic closure of  $k$  such that the absolute Galois group  $\text{Gal}(\bar{k}=k)$  of  $k$  with respect to  $\bar{k}$  is naturally identified with the quotient  $G_k/I_k$ . We shall write

$\text{Frob}_k \in \text{Gal}(\bar{k}=k) \setminus G_k/I_k$  for the ( $|k|$ -th power) Frobenius element.

We shall write, moreover,

$\text{Br}(k) \stackrel{\text{def}}{=} H^2(G_k; \bar{k})$  (where we regard  $\bar{k}$  as a discrete  $G_k$ -module) for the Brauer group of  $k$ .

The following lemma is one of fundamental results concerning the structure of the topological group  $G_k$ .

Lemma 1.4 . The following hold:

- (i) The topological group  $G_k$  is topologically finitely generated.
- (ii) For a subgroup  $H \subset G_k$  of  $G_k$ , it holds that  $H$  is open in  $G_k$  if and only if  $H$  is of finite index in  $G_k$ .

Proof. Assertion (i) follows from [9, Theorem 7.4.1]. Assertion (ii) follows from [10, Theorem 1.1], together with assertion (i).

The following lemma is one of fundamental results concerning the structure of the tame quotient  $G_k/P_k$  of  $G_k$ .

Lemma 1.5 . The following hold:

- (i) The quotient  $G_k/I_k$  is topologically generated by  $\text{Frob}_k \in G_k/I_k$  and, as an abstract topological group, isomorphic to  $\mathbb{Z}_+$ . In particular, the quotient  $G_k/I_k$  is abelian.

(ii) There exists an isomorphism of topological groups

$$I_k = P_k \cong (\bar{k})^{(p_k^0)};$$

where we write  $(\bar{k})^{(p_k^0)}$  for the quotient of  $(\bar{k})$  by the pro- $p_k$ -Sylow subgroup of  $(\bar{k})$ . Moreover, such an isomorphism is always  $G_k$ -equivariant (i.e., with respect to the action of  $G_k$  on  $I_k = P_k$  by conjugation and the natural action of  $G_k$  on  $(\bar{k})^{(p_k^0)}$ ).

(iii) The action of  $G_k$  on  $I_k = P_k$  by conjugation determines an injection

$$G_k = I_k \rightarrow \prod_{l: \text{prime}; l \nmid p_k} Z_l = \text{Aut}(I_k = P_k)$$

(cf. (ii)) which maps  $\text{Frob}_k \in G_k = I_k$  to  $p_k^{f_k} \in \prod_{l: \text{prime}; l \nmid p_k} Z_l$ .

Proof. Assertion (i) is discussed in the discussion following [9, Proposition 7.5.1]. Assertion (ii) follows immediately from [9, Proposition 7.5.2]. Assertion (iii) follows immediately in light of assertion (ii) from [9, Lemma 7.5.4(ii)], together with the definition of  $\text{Frob}_k \in G_k = I_k$ .

Next, let us recall an explicit description of the Brauer group of an MLF as follows.

Lemma 1.6 . Write  $k_{ur} = \bar{k}$  for the maximal unramified extension of  $k$  in  $\bar{k}$  (i.e.,  $k_{ur} = \bar{k}^{I_k}$ ),  $O_{k_{ur}} = O_k$  for the ring of integers of  $k_{ur}$ , and  $V \stackrel{\text{def}}{=} k_{ur} = O_{k_{ur}}$ . Then the following hold:

(i) The monoids  $O_{k_{ur}}^B = O_{k_{ur}} \rtimes V$  are, as abstract monoids, isomorphic to  $N \rtimes Z_+$ . Moreover, the action of  $G_k$  on  $V$  determined by the action of  $G_k$  on  $k_{ur}$  is trivial.

(ii) For each positive integer  $n$ , the natural homomorphism

$$H^2(G_k; (\bar{k})[n]) \rightarrow \text{Br}(k)$$

determines an isomorphism

$$H^2(G_k; (\bar{k})[n]) \cong \text{Br}(k)[n];$$

(iii) The natural homomorphism

$$H^2(G_k = I_k; k_{ur}) \cong \text{Br}(k)$$

is an isomorphism

(iv) The natural homomorphism

$$H^2(G_k = I_k; k_{ur}) \cong H^2(G_k = I_k; V)$$

is an isomorphism

(v) The homomorphism

$$H^1(G_k = I_k; V^{pf} = V) \cong H^2(G_k = I_k; V)$$

determined by the exact sequence of  $G_k = I_k$ -modules

$$0 \rightarrow V \rightarrow V^{pf} \rightarrow V^{pf} = V \rightarrow 0$$

is an isomorphism

(vi) The homomorphism

$$H^1(G_k = I_k; V^{pf} = V) \rightarrow V^{pf} = V$$

obtained by mapping

$$2 \text{ Hom}(G_k = I_k; V^{pf} = V) = H^1(G_k = I_k; V^{pf} = V)$$

(cf. (i)) to

$$(\text{Frob}_k) \otimes V^{pf} = V$$

is an isomorphism

(vii) The various isomorphisms of (iii), (iv), (v), (vi) determine an isomorphism

$$\text{Br}(k) \rightarrow V^{pf} = V$$

(which thus determines an isomorphism

$$H^2(G_k; (\bar{k})) \rightarrow V^{\wedge}$$

cf. (i), (ii)).

Proof. Assertion (i) follows immediately from the (easily verified) fact that the natural inclusion  $k \rightarrow k_{ur}$  determines isomorphisms  $O_k^B = O_k \rightarrow O_{k_{ur}}^B = O_{k_{ur}}, k = O_k \rightarrow V$ . Assertion (ii) follows immediately from Hilbert Theorem 90 (cf. [9, Theorem 6.2.1]). Assertion (iii) follows from [11, §1.1, Theorem 1]. Assertion (iv) follows from [11, §1.1, Theorem 2]. Assertion (v) follows from the discussion following [11, §1.1, Theorem 2]. Assertion (vi) follows from the discussion preceding [11, §1.1, Corollary]. Assertion (vii) follows from [11, §1.1, Corollary].

Next, let us recall local class field theory as follows.

Lemma 1.7 . There exists an injective homomorphism

$$\text{req}_k : k \rightarrow G_k^{ab}$$

which satisfies the following conditions:

(i) The homomorphism  $\text{req}_k$  determines a commutative diagram

$$\begin{array}{ccccccc}
 1 & \rightarrow & O & \xrightarrow{\gamma^k} & k & \rightarrow & k = O_k & \rightarrow & 1 \\
 & & & \uparrow \text{?} & \uparrow \text{?} & & \uparrow \text{?} & & \\
 & & & \alpha^y & \alpha^y & & \alpha^y & & \\
 1 & \rightarrow & \text{Im}(I_k \rightarrow G_k) & \rightarrow & G_k^{ab} & \rightarrow & G_k^{ab} & \xrightarrow{\text{Frob}_k^Z} & 1 \\
 & & & & & & \uparrow \text{?} & & \\
 & & & & & & \alpha^y & & \\
 1 & \rightarrow & \text{Im}(I_k \rightarrow G_k) & \rightarrow & G_k^{ab} & \rightarrow & G_k = I_k & \rightarrow & 1
 \end{array}$$

(cf. Lemma 1.5(i)) where the horizontal sequences are exact, the upper vertical arrows are isomorphisms, the lower vertical arrows are injective, and the right-hand upper vertical arrow  $k = O_k \rightarrow \text{Frob}_k^Z$  determines an isomorphism

$$O_k^B = O_k \rightarrow \text{Frob}_k^N:$$

In particular, the homomorphism  $\text{req}_k$  determines an isomorphism of topological modules

$$(k^{\wedge}) \rightarrow G_k^{ab}:$$

(ii) Let  $K/\bar{k}$  be a finite extension of  $k$ . (So  $K$  is an MLF, and  $G_K \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=K) \leq G_k$  is an open subgroup of  $G_k$ .) Then the Norm map  $\text{Nm}_{K=k} : K \rightarrow k$  and the natural homomorphism  $G_K^{\text{ab}} \rightarrow G_k^{\text{ab}}$  fit into the following commutative diagram:

$$\begin{array}{ccc} K & \xrightarrow{\text{Nm}_{K=k}} & k \\ \text{rec}_K \downarrow \cong & & \text{rec}_k \downarrow \cong \\ G_K^{\text{ab}} & \xrightarrow{\quad} & G_k^{\text{ab}} \end{array}$$

(iii) Let  $K/\bar{k}$  be a finite extension of  $k$ . (So  $K$  is an MLF, and  $G_K \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=K) \leq G_k$  is an open subgroup of  $G_k$ .) Then the natural inclusion  $k \hookrightarrow K$  and the transfer map  $\text{Tf}_{G_K/G_k} : G_K^{\text{ab}} \rightarrow G_k^{\text{ab}}$  fit into the following commutative diagram:

$$\begin{array}{ccc} k & \xrightarrow{\quad} & K \\ \text{rec}_k \downarrow \cong & & \text{rec}_K \downarrow \cong \\ G_k^{\text{ab}} & \xrightarrow{\text{Tf}_{G_K/G_k}} & G_K^{\text{ab}} \end{array}$$

(iv) Let  $L$  be an MLF,  $\bar{L}$  an algebraic closure of  $L$ , and  $\sigma : k \xrightarrow{\cong} L$  an isomorphism of fields. Then the diagram

$$\begin{array}{ccc} k & \xrightarrow{\quad} & L \\ \text{rec}_k \downarrow \cong & & \text{rec}_L \downarrow \cong \\ G_k^{\text{ab}} & \xrightarrow{\quad} & \text{Gal}(\bar{L}=L)^{\text{ab}} \end{array}$$

(where the lower horizontal arrow is the isomorphism induced by  $\sigma$ ) commutes

Proof. This assertion follows immediately from the various assertions in [11, §2].

Finally, as an application of local class field theory, let us verify the slimness of the absolute Galois group of an MLF.

Lemma 1.8. The absolute Galois group  $G_k$  is slim, hence also center-free

Proof. Let  $K$  be a finite extension of  $k$  contained in  $\bar{k}$  and  $\alpha \in G_k$  an element of the centralizer of  $G_K \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=K)$  in  $G_k$ . Let us observe that, to verify  $\alpha = 1$ , we may assume without loss of generality, by replacing  $K$  by a suitable finite extension of  $K$  contained in  $\bar{k}$ , that  $K$  is Galois over  $k$ . Thus, it follows immediately from the injectivity of the homomorphism  $\text{rec}_K$  of Lemma 1.7, together with Lemma 1.7 (iv), that the action of  $G_k/G_K = \text{Gal}(K=k)$  on  $G_K^{\text{ab}}$  by conjugation is faithful, which thus implies that  $\alpha \notin G_K$ . In particular, by allowing  $K$  to vary, we conclude that  $\alpha = 1$ , as desired. This completes the proof of Lemma 1.8.

## 2. Bi-anabelian Results for MLF's

In the present section, let us recall some results in bi-anabelian geometry for MLF's. In the present section, for  $f \neq g$ , let  $k$  be an MLF and  $\bar{k}$  an algebraic closure of  $k$ ; write

$G \stackrel{\text{def}}{=} \text{Gal}(\bar{k} = k)$ . Thus, we have a natural map

$$= \text{Isom}(k ; k) \rightarrow \text{Isom}(G ; G) = \text{Inn}(G) :$$

Typically, bi-anabelian geometry (i.e., the classical point of view of anabelian geometry) discusses some properties such as faithfulness/ fullness of the (restriction, to a certain suitable category of geometric objects, of the) functor of taking arithmetic fundamental groups. Put another way, in a discussion of bi-anabelian geometry, one usually sees two schemes/rings of interest (e.g., the two fields  $k$  and  $\bar{k}$  in our case) and discusses the relationship between a certain set of morphisms (e.g., the set of isomorphisms) between the fixed two schemes/rings and a certain set of homomorphisms (e.g., the set of isomorphisms) between the arithmetic fundamental groups. In particular, roughly speaking, bi-anabelian geometry (i.e., for isomorphisms between MLF's) may be summarized as the study of the above map.

The following proposition asserts the injectivity of the map .

Proposition 2.1 . The map is injective.

Proof. Since (one verifies easily that) every automorphism of the field  $\bar{k}$  is an automorphism over  $\mathbb{Q}_{p_k}$  (  $\bar{k}$  ), this assertion follows immediately, by considering the difference of two elements of  $\text{Isom}(k ; k)$  whose images via coincide, from Lemma 1.8.

The following theorem is fundamental in bi-anabelian geometry for MLF's.

Theorem 2.2 . The following hold:

- (i) There exists a pair  $(\bar{k} = k ; \bar{k} = k)$  which satisfies the following condition: The domain of  $\text{Isom}(k ; k)$  is empty, but the codomain of  $\text{Isom}(G ; G)$  is nonempty.
- (ii) There exists a pair  $(\bar{k} = k ; \bar{k} = k)$  which satisfies the following condition: The domain of  $\text{Isom}(k ; k)$  is nonempty (which thus implies that the codomain of  $\text{Isom}(G ; G)$  is nonempty), but the map  $\text{Isom}(k ; k) \rightarrow \text{Isom}(G ; G)$  is not surjective.

Proof. Assertion (i) follows from the examples discussed in [13, §2]. Assertion (ii) follows from the discussion given at the final portion of [9, Chapter VII, §5].

Remark 2.2.1 . In [3], a necessary and sufficient condition for the pair  $(\bar{k} = k ; \bar{k} = k)$  to satisfy that the codomain of  $\text{Isom}(G ; G)$  is nonempty was discussed.

The map is always injective (cf. Proposition 2.1) but not surjective in general (cf. Theorem 2.2). Thus, in the study of bi-anabelian geometry for MLF's, one often discusses conditions for an outer isomorphism  $G \rightarrow G$  (i.e., an element of the codomain of ) to be contained in the image of . The following theorem gives such conditions.

Theorem 2.3 . Let  $\phi : G \rightarrow G$  be an isomorphism of topological groups (which thus implies that  $p_k = p_k$  cf. Remark 2.3.1 below). Then the following two conditions are equivalent:

- (i) The outer isomorphism determined by  $\phi$  (i.e., the element of the codomain of  $\text{Isom}(G ; G)$  determined by  $\phi$ ) is contained in the image of  $\text{Isom}(k ; k) \rightarrow \text{Isom}(G ; G)$ .

- (ii) The isomorphism  $\alpha$  is compatible with the respective ramification filtrations (cf. [11, §4.1]) of  $G, G$ .
- (iii) For  $2 \leq f \leq g$ , write  $(\bar{R})_+$  for the topological  $G$ -module obtained by forming the underlying additive module of the  $p_k$ -adic completion  $\bar{R}$  of  $\bar{k}$ . Write, moreover,  $(\bar{R})_+$  for the topological  $G$ -module obtained by considering the action of  $G$  on  $(\bar{R})_+$  via  $\alpha$ . Then there exists a  $G$ -equivariant topological isomorphism  $(\bar{R})_+ \xrightarrow{\sim} (\bar{R})_+$ .
- (iv) Write  $(O_k)_+$  for the  $G$ -module obtained by considering the action of  $G$  on  $(O_k)_+$  via  $\alpha$ . Then there exists a  $G$ -equivariant isomorphism  $(O_k)_+ \xrightarrow{\sim} (O_k)_+$ .
- (v) For every finite-dimensional Hodge Tate representation  $\rho$  of  $G$  over  $\mathbb{Q}_{p_k} = \mathbb{Q}_{p_k}$ , the finite-dimensional representation of  $G$  over  $\mathbb{Q}_{p_k} = \mathbb{Q}_{p_k}$  obtained by forming the composite  $\rho \circ \alpha$  is Hodge Tate.

*Proof.* The equivalence (i), (ii) follows from [4, Theorem]. The equivalence (i), (iii) follows from [5, Theorem 3.5(ii)]. The implications (i)  $\Rightarrow$  (iv)  $\Rightarrow$  (iii) are immediate; thus, by the equivalence (i), (iii) already discussed, we obtain the equivalence (i), (iv). The equivalence (i), (v) follows from [1, Theorem].

**Remark 2.3.1.** Suppose that the codomain of  $\alpha$  is nonempty, i.e., that there exists an isomorphism  $\alpha : G \xrightarrow{\sim} G$  of topological groups. Then it is well-known that it holds that  $p_k = p_k$ . This fact also follows from Proposition 3.6 of the present article.

**Remark 2.3.2.**

- (i) One may find other conditions for  $\alpha$  as in Theorem 2.3 equivalent to condition (i) of Theorem 2.3 in, for instance, [5, §3], [1, §3].
- (ii) We have considered, in Theorem 2.3, some conditions for another isomorphism between the absolute Galois groups of MLF's to arise from an isomorphism between the original MLF's. On the other hand, one may consider a condition for another open homomorphism between the absolute Galois groups of MLF's to arise from an homomorphism between the original MLF's. One may also find such conditions in, for instance, [5, §3], [1, §3]. For instance, a similar equivalence to the equivalence (i)  $\Rightarrow$  (v) of Theorem 2.3 still holds even if one considers an open homomorphism (i.e., as opposed to an isomorphism) from  $G$  to  $G$  (cf. [1, Theorem]).

### 3. Mono-anabelian Reconstruction for MLF's: I

Let us recall that, as discussed at the beginning of Section 2, anabelian geometry centers around a comparison between two fixed schemes/rings of interest via the arithmetic fundamental groups. By contrast, mono-anabelian geometry centers around the task of establishing a group-theoretic software (i.e., group-theoretic algorithm) whose input data consists of a single abstract (topological) group isomorphic to the arithmetic fundamental

group of a scheme/ring of interest (cf., e.g., [6, Introduction, Remark 1.9.8, and Remarks following Corollary 3.7] for more details concerning bi-anabelian/mono-anabelian geometry).

$$\begin{array}{l}
 \text{bi-anabelian geometry} \\
 \text{et}_1(X) \cong \text{et}_1(X) \Rightarrow \text{(objects related to } X \text{)} \cong \text{(objects related to } X \text{)} \\
 \text{mono-anabelian geometry} \\
 \text{an isomorph of } \text{et}_1(X) \cong \text{isomorph(s) of (objects related to } X \text{)}
 \end{array}$$

In the present section, let us establish some mono-anabelian reconstruction algorithms for MLF's. In particular, we discuss some group-theoretic algorithms (cf. Remark 3.15.1 below) whose input data consist of an abstract group isomorphic to the absolute Galois group of an MLF.

**Definition 3.1** . We shall refer to an isomorph, as a group, of the absolute Galois group of an MLF as a group of MLF-type.

In the remainder of the present section, let  $G$  be a group of MLF-type,  $k$  an MLF, and  $\bar{k}$  an algebraic closure of  $k$ . We shall also apply the notational conventions related to  $k$  and  $\bar{k}$  introduced in Section 1.

**Definition 3.2** . We shall say that a subgroup of  $G$  is open if the subgroup is of finite index in  $G$ .

**Proposition 3.3** . The following hold:

- (i) The open subgroups of  $G$  (i.e., in the sense of Definition 3.2) determine a structure of profinite group of  $G$ .
- (ii) Every isomorphism  $G \cong G_k$  of groups is an isomorphism of topological groups with respect to the structure of profinite group of  $G$  of (i).

**Proof.** Assertion (i) follows from Lemma 1.4(ii). Assertion (ii) follows from (i).

In the remainder of the present article, we always regard a group of MLF-type as a profinite, hence also topological, group by Proposition 3.3(i). Note that it follows from the various definitions involved that every open subgroup of a group of MLF-type is of MLF-type.

**Lemma 3.4** . The following hold:

- (i) There exists a uniquely determined prime number  $l$  such that

$$\log_l |(G^{ab=tor}) / G^{ab=tor}| \leq 2;$$

- (ii) The subquotient  $G^{ab=tor}$  of  $G$  is finite .

**Proof.** Let us observe that it follows from Lemma 1.7(i), together with Lemma 1.2(i), that the topological module  $G_k^{ab}$  is isomorphic to the topological module

$$Z_+ = (p_k^f - 1)Z_+ \quad (Z_+ = p_k^a Z_+) \quad (Z_{p_k})_+^{d_k} \oplus \mathbb{Z}_+$$



for some nonnegative integers. In particular, the topological module  $G_k^{ab=tor}$  (respectively,  $G_k^{ab-tor}$ ) is isomorphic to the topological module

$$(Z_{p_k})_+^{d_k} \oplus \mathbb{Z}_+ \quad (\text{respectively, } Z_+ = (p_k^{f_k} - 1)Z_+ \quad (Z_+ = p_k^a Z_+)):$$

Thus, assertion (i) (respectively, (ii)) holds.

Definition 3.5 .

(i) We shall write

$$p(G)$$

for the uniquely determined (cf. Lemma 3.4(i)) prime number such that

$$\log_{p(G)} ] G^{ab=tor} = p(G) G^{ab=tor} \quad 2:$$

(ii) We shall write

$$d(G) \stackrel{\text{def}}{=} \log_{p(G)} ] G^{ab=tor} = p(G) G^{ab=tor} \quad 1;$$

$$f(G) \stackrel{\text{def}}{=} \log_{p(G)} 1 + ] (G^{ab-tor})^{(p(G)^0)} \quad (\neq 0);$$

where we write  $(G^{ab-tor})^{(p(G)^0)}$  for the quotient of  $G^{ab-tor}$  by the  $p(G)$ -Sylow subgroup of (the finite cf. Lemma 3.4(ii) abelian group)  $G^{ab-tor}$ , and

$$e(G) \stackrel{\text{def}}{=} d(G) = f(G):$$

(iii) We shall write

$$I(G) \stackrel{\text{def}}{=} \bigcup_{N \triangleleft G} N;$$

where  $N$  ranges over the normal open subgroups  $\mathfrak{O}\mathfrak{G}$  (so  $N$  is of MLF-type cf. the discussion following Proposition 3.3) such that  $e(N) = e(G)$ ;

$$P(G) \stackrel{\text{def}}{=} \bigcup_{N \triangleleft G} N;$$

where  $N$  ranges over the normal open subgroups  $\mathfrak{O}\mathfrak{G}$  (so  $N$  is of MLF-type cf. the discussion following Proposition 3.3) such that  $e(N) = e(G)$  is a positive integer prime to  $p(G)$ .

Proposition 3.6 . It holds that

$$p_k = p(G_k); \quad d_k = d(G_k); \quad f_k = f(G_k); \quad e_k = e(G_k);$$

$$I_k = I(G_k); \quad P_k = P(G_k):$$

Proof. The assertions for  $p_k$ ,  $d_k$ , and  $f_k$  follow from Lemma 1.7(i), together with Lemma 1.2(i) (cf. also the explicit descriptions of  $G_k^{ab=tor}$  and  $G_k^{ab-tor}$  given in the proof of Lemma 3.4). The assertion for  $e_k$  follows in light of the assertions for  $d_k$  and  $f_k$  from Lemma 1.2(iii). The assertions for  $I_k$  and  $P_k$  follow in light of the assertions for  $p_k$  and  $e_k$  from the definitions of  $I_k$  and  $P_k$ .

Lemma 3.7 . The following hold:

- (i) The quotients  $G/I(G)$  and  $I(G)=P(G)$  are abelian.
- (ii) There exists a uniquely determined element  $z \in G/I(G)$  of  $G/I(G)$  such that the action of  $z$  on (the abelian cf. (i) group)  $I(G)=P(G)$  by conjugation is given by multiplication by  $p(G)^{f(G)}$ .

Proof. Assertion (i) follows in light of Proposition 3.6 from Lemma 1.5(i), (ii). Assertion (ii) follows in light of Proposition 3.6 from Lemma 1.5(iii).

Definition 3.8 . We shall write

$$\text{Frob}(G) \in G/I(G)$$

for the uniquely determined (cf. Lemma 3.7(ii)) element of  $G/I(G)$  such that the action of  $\text{Frob}(G)$  on  $I(G)=P(G)$  by conjugation is given by multiplication by  $p(G)^{f(G)}$  (cf. Lemma 3.7(i)).

Proposition 3.9 . It holds that

$$\text{Frob}_k = \text{Frob}(G_k)$$

in  $G_k/I_k = G_k/I(G_k)$  (cf. Proposition 3.6).

Proof. This assertion follows in light of Proposition 3.6 from Lemma 1.5(iii).

Definition 3.10 .

(i) We shall write

$$O(G) \stackrel{\text{def}}{=} \text{Im } I(G) \rightarrow G \rightarrow G^{\text{ab}} \rightarrow G^{\text{ab}}.$$

By considering the topology induced by the topology of  $I(G)$ , we regard  $O(G)$  as a pro finite, hence also topological, module.

(ii) We shall write

$$O(G) \supseteq O(G)$$

for the maximal pro- $p(G)$  submodule of the pro finite module  $O(G)$ .

(iii) We shall write

$$\underline{k}(G) \stackrel{\text{def}}{=} O(G) = O(G):$$

(iv) We shall write

$$k(G) \stackrel{\text{def}}{=} G^{\text{ab}} \times_{G/I(G)} \text{Frob}(G)^{\mathbb{Z}} \times G^{\text{ab}}$$

(cf. Lemma 3.7(i)),

$$\text{rec}(G): k(G) \rightarrow G^{\text{ab}}$$

for the natural inclusion, and

$$O^B(G) \stackrel{\text{def}}{=} G^{\text{ab}} \times_{G/I(G)} \text{Frob}(G)^{\mathbb{N}} \times k(G)$$

(cf. Lemma 3.7(i)). By considering the topologies induced by the topology of  $O(G)$  (cf (i)), we regard  $O^B(G)$ ,  $k(G)$  as a topological monoid, topological module, respectively.

(v) We shall write

$$k(G) \stackrel{\text{def}}{=} k(G)^\sim$$

and

$$\underline{k}(G) \stackrel{\text{def}}{=} \underline{k}(G)^\sim :$$

(vi) We shall write

$$k_+(G) \stackrel{\text{def}}{=} O(G)^{\text{pf}}$$

and

$$I(G) \stackrel{\text{def}}{=} 2 \cdot \rho(G)^{-1} \text{Im } O(G) \rightarrow k_+(G) \rightarrow k_+(G) :$$

By considering the topologies induced by the topology of  $O(G)$  (cf. (i)), we regard  $k_+(G), I(G)$  as topological modules, respectively.

Proposition 3.11 . The following hold:

(i) The injective homomorphism  $\text{rec}_k : k \rightarrow G_k^{\text{ab}}$  of Lemma 1.7 determines a commutative diagram of topological monoids

$$\begin{array}{ccccccc} O_k & \rightarrow & O & \rightarrow & O^B & \rightarrow & k \rightarrow \text{rec}_k \rightarrow G_k^{\text{ab}} \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ O(G_k) & \rightarrow & O(G_k) & \rightarrow & O^B(G_k) & \rightarrow & k(G_k) \rightarrow \text{rec}(G_k) \rightarrow G_k^{\text{ab}} \end{array}$$

where the horizontal arrows are injective, and the vertical arrows are isomorphisms

(ii) The isomorphism  $k \rightarrow k(G_k)$  in the diagram of (i) determines an isomorphism of monoids

$$k \rightarrow k(G_k) :$$

(iii) The left-hand square of the diagram of (i) determines isomorphisms of monoids

$$\underline{k} \rightarrow \underline{k}(G_k); \quad \underline{k} \rightarrow \underline{k}(G_k) :$$

(iv) The isomorphism  $O_k \rightarrow O(G_k)$  in the diagram of (i) and the natural homomorphisms  $O(G_k) \rightarrow I(G_k) \rightarrow k_+(G_k)$  fit into a commutative diagram of topological modules

$$\begin{array}{ccc} O_k & \rightarrow & \text{log}_k \rightarrow \\ \downarrow \cong & & \downarrow \cong \\ O(G_k) & \rightarrow & I(G_k) \rightarrow k_+(G_k) \end{array}$$

where the vertical arrows are isomorphisms

Proof. Assertion (i) follows immediately in light of Proposition 3.6 and Proposition 3.9 from Lemma 1.7(i), together with Lemma 1.2(ii). Assertions (ii), (iii) are immediate. Assertion (iv) follows in light of Proposition 3.6 from Lemma 1.2(iv).

Lemma 3.12 . The following hold:

(i) The topological module  $k_+(G)$  is locally compact.

- (ii) The topological submodule  $I(G) \subseteq k_+(G)$  of  $k_+(G)$  is compact and open.
- (iii) For each positive real number  $a$ , there exists a uniquely determined Haar measure on  $k_+(G)$  (cf. (i)) that assigns the compact open (cf. (ii)) subset  $I(G) \subseteq k_+(G)$  to  $a$ .

Proof. Assertions (i), (ii) follow from Proposition 3.11 (iv). Assertion (iii) follows from assertions (i), (ii).

Definition 3.13 . We shall write

$$p(G)$$

for the uniquely determined (cf. Lemma 3.12 (iii)) Haar measure on  $k_+(G)$  such that

$$p(G) (I(G)) = p(G)^{d(G) + f(G)} |k(G)_{\text{tor}}|^{p(G)};$$

where we write

$$p(G) \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } p(G) \neq 2 \\ 2 & \text{if } p(G) = 2 \end{cases}$$

and  $|k(G)_{\text{tor}}|^{p(G)}$  for the  $p(G)$ -Sylow subgroup of (the finite abelian group)  $k(G)_{\text{tor}}$ . (cf. Lemma 1.2 (i), Proposition 3.11 (i))

Proposition 3.14 . It holds that

$$k = \varprojlim_k (G_k)$$

relative to the right-hand vertical arrow  $k_+ \rightarrow k_+(G_k)$  of the diagram of Proposition 3.11 (iv).

Proof. This assertion follows in light of Proposition 3.6 and Proposition 3.11 (i), (iv) from Lemma 1.3 (iii).

The various assertions discussed in the present section may be summarized as follows.

Summary 3.15 . There exist functorial group-theoretic algorithms (cf. Remark 3.15.1 below) for constructing, from a group  $G$  of MLF-type,

- a topology on  $G$ ,
- a prime number  $p(G)$ ,
- positive integers  $d(G)$ ,  $f(G)$ , and  $e(G)$ ,
- subgroups  $P(G) \subseteq I(G) \subseteq G$  of  $G$ ,
- an element  $\text{Frob}(G) \in G = I(G)$  of  $G = I(G)$ ,
- topological monoids  $O(G) \subseteq O(G) \subseteq O^B(G) \subseteq k(G) \xrightarrow{\text{rec}(G)} G^{\text{ab}}$ ,
- monoids  $\underline{k}(G) \subseteq \underline{k}(G)$  and  $k(G)$ ,
- topological modules  $I(G) \subseteq k_+(G)$ , and
- a measure  $p(G)$  on  $k_+(G)$

which correspond to

the pro finite topology on  $G_k$ ,

the prime number  $p_k$ ,

the positive integers  $d_k$ ,  $f_k$ , and  $e_k$ ,

the subgroups  $P_k = I_k = G_k$  of  $G_k$ ,

the element  $\text{Frob}_k \in G_k = I_k$  of  $G_k = I_k$ ,

the topological monoids  $O_k = O_k = O_k^B = k \prod_r^{\text{rec}_k} G_k^{\text{ab}}$ ,

the monoids  $\underline{k} = \underline{k}$  and  $k_+$ ,

the topological modules  $\underline{k} = k_+$ , and

the measure  $\mu_k$  on  $k_+$ ,

respectively.

**Remark 3.15.1** . As discussed in [6, Remark 1.9.8], in anabelian geometry the term group-theoretic is usually used simply to mean preserved by an arbitrary isomorphism between the arithmetic fundamental groups under consideration. On the other hand, in mono-anabelian geometry the term group-theoretic algorithm is used to mean that the algorithm in a discussion is phrased in language that only depends on the topological group structure of the arithmetic fundamental group under consideration.

**Remark 3.15.2** .

(i) It follows from Theorem 2.2(ii) and the equivalence (i), (ii) of Theorem 2.3 that there exist an MLF  $L$ , an algebraic closure  $\bar{L}$  of  $L$ , and an automorphism  $\sigma$  of  $G_L \stackrel{\text{def}}{=} \text{Gal}(\bar{L}=L)$  not compatible with the ramification filtration of  $G_L$ . By this fact, one may conclude that the ramification filtration of the absolute Galois group of an MLF should be considered to be not group-theoretic.

(ii) It follows from Theorem 2.2(ii) and the equivalence (i), (v) of Theorem 2.3 that there exist an MLF  $L$ , an algebraic closure  $\bar{L}$  of  $L$ , an automorphism  $\sigma$  of  $G_L \stackrel{\text{def}}{=} \text{Gal}(\bar{L}=L)$ , and a finite-dimensional Hodge Tate representation  $\rho$  of  $G_L$  over  $\mathbb{Q}_p$  such that the composite  $\rho \circ \sigma$  is not Hodge Tate (cf. also [1, Remark 3.3.1]). By this fact, one may conclude that Hodge Tate-ness of the  $p$ -adic representations of the absolute Galois group of an MLF should be considered to be not group-theoretic.

#### 4. Mono-anabelian Reconstruction for MLF's: II

In the present section, we continue to establish mono-anabelian reconstruction algorithms for MLF's. Let  $G$  be a group of MLF-type,  $k$  an MLF, and  $\bar{k}$  an algebraic closure of  $k$ . We shall also apply the notational conventions related to  $k$  and  $\bar{k}$  introduced in Section 1.

Definition 4.1 .

(i) We shall write

$$\begin{aligned} \bar{O}(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} O(H); & \bar{k}(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} k(H); & \bar{k}(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} k(H); \\ \bar{O}^B(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} O^B(H); & \bar{k}(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} k(H); & \bar{k}(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} k(H); \\ & & \bar{k}_+(G) &\stackrel{\text{def}}{=} \varinjlim_{H \subset G} k_+(H); \end{aligned}$$

where the inductive limits are taken over the open subgroups  $H \subset G$  of  $G$  (so  $H$  is of MLF-type cf. the discussion following Proposition 3.3), and the transition morphisms in the limits are given by the homomorphisms determined by the transfer maps. We regard these seven monoids as  $G$ -monoids by the actions of  $G$  obtained by conjugation.

(ii) We shall write

$$(G) \stackrel{\text{def}}{=} \bar{k}(G)_{\text{tor}} :$$

We regard  $(G)$  as a  $G$ -module by the action of  $G$  induced by the action of  $G$  on  $\bar{k}(G)$  (cf. (i)).

(iii) We shall write

$$(G) \stackrel{\text{def}}{=} \varprojlim_n (G)[n]$$

where the projective limit is taken over the positive integers  $n$  and refer to  $(G)$  as the cyclotome associated to  $G$ . We regard the cyclotome  $(G)$  as a pro finite, hence also topological, module (by the easily verified finiteness of  $(G)[n]$  for each  $n$ ). Moreover, we also regard the cyclotome  $(G)$  as a (topological)  $G$ -module by the action of  $G$  induced by the action of  $G$  on  $(G)$  (cf. (ii)).

Proposition 4.2 . The following hold:

(i) The various injective homomorphisms  $\text{res}_K : K \rightarrow G_K^{\text{ab}}$  where  $K$  ranges over the finite extensions of  $k$  contained in  $\bar{k}$  of Lemma 1.7 determine a commutative diagram of  $G_k$ -monoids

$$\begin{array}{ccccccc} O_k & \hookrightarrow & O & \xrightarrow{B} & \bar{k} & \hookrightarrow & \bar{k} \\ \uparrow \cong & & \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\ \bar{O}(G_k) & \hookrightarrow & \bar{O}^B(G_k) & \hookrightarrow & \bar{k}(G_k) & \hookrightarrow & \bar{k}(G_k) \end{array}$$

where the horizontal arrows are injective, and the vertical arrows are isomorphisms

(ii) The isomorphism  $O_k \xrightarrow{\cong} \bar{O}(G_k)$  in the diagram of (i) determines isomorphisms of  $G_k$ -monoids

$$\bar{k} \xrightarrow{\cong} \bar{k}(G_k); \quad \bar{k} \xrightarrow{\cong} \bar{k}(G_k):$$

(iii) The isomorphism  $O_k \cong \bar{O}(G_k)$  in the diagram of (i) and the natural homomorphism  $\bar{O}(G_k) \rightarrow \bar{k}_+(G_k)$  fit into a commutative diagram of  $G_k$ -modules

$$\begin{array}{ccc} O_k & \xrightarrow{\log_k} & \bar{k}_+ \\ \downarrow \cong & & \downarrow \cong \\ \bar{O}(G_k) & \longrightarrow & \bar{k}_+(G_k) \end{array}$$

where the vertical arrows are isomorphisms

(iv) The isomorphism  $\bar{k} \cong \bar{k}(G_k)$  in the diagram of (i) determines an isomorphism of  $G_k$ -modules

$$\bar{k} \cong \bar{k}(G_k);$$

hence also an isomorphism of topological  $G_k$ -modules

$$(\bar{k}) \cong (\bar{k}(G_k));$$

Proof. These assertions follow in light of Proposition 3.11 (i), (ii), (iii), (iv) from Lemma 1.7 (iii), (iv).

The various assertions discussed in the present section may be summarized as follows.

Summary 4.3 . There exist functorial group-theoretic algorithms (cf. Remark 3.15.1) for constructing, from a group  $G$  of MLF-type,

$G$ -monoids  $\bar{O}(G) \cong \bar{O}^B(G) \cong \bar{k}(G) \cong \bar{k}(G)$ ,

$G$ -monoids  $\bar{k}(G) \cong \bar{k}(G)$ ,

a  $G$ -module  $\bar{k}_+(G)$ ,

a  $G$ -module  $(\bar{k})$ , and

a topological  $G$ -module  $(\bar{k})$

which correspond to

the  $G_k$ -monoids  $O_k \cong O_k^B \cong \bar{k} \cong \bar{k}$ ,

the  $G_k$ -monoids  $\bar{k} \cong \bar{k}$ ,

the  $G_k$ -module  $\bar{k}_+$ ,

the  $G_k$ -module  $(\bar{k})$ , and

the topological  $G_k$ -module  $(\bar{k})$ ,

respectively.

Remark 4.3.1 .

(i) It follows from Theorem 2.2(ii) and the equivalence (i) , (iii) of Theorem 2.3 that there exist an MLF  $L$ , an algebraic closure  $\bar{L}$  of  $L$ , and an automorphism  $\sigma$  of  $\text{Gal}(\bar{L}=L)$  such that the automorphism of  $\bar{L}_+$  induced by  $\sigma$  (cf. Proposition 4.2(iii)) is not compatible with the  $p_L$ -adic topology of  $\bar{L}_+$ . By this fact, one may conclude that

the  $p$ -adic topology on the underlying additive module of an algebraic closure of an MLF should be considered to be not group-theoretic

(cf. also Remark 4.3.2(ii) below).

(ii) It follows from Theorem 2.2(ii) and the equivalence (i) , (iv) of Theorem 2.3 that there exist an MLF  $L$ , an algebraic closure  $\bar{L}$  of  $L$ , and an automorphism  $\sigma$  of  $\text{Gal}(\bar{L}=L)$  such that the automorphism of  $\bar{L}_+$  induced by  $\sigma$  (cf. Proposition 4.2(iii)) does not restrict to an automorphism of the submodule  $(O_L)_+ \subset \bar{L}_+$  of  $\bar{L}_+$ . By this fact, one may conclude that

the underlying additive module of the ring of integers of an algebraic closure of an MLF should be considered to be not group-theoretic .

(iii) Assume that there exists a functorial group-theoretic algorithm for constructing, from a group  $G$  of MLF-type, the submodule of  $k_+(G)$  which corresponds to the submodule  $(O_k)_+ \subset k_+$  of  $k_+$ . Then, by applying a similar construction to the constructions of Definition 4.1 (i), one may obtain a functorial group-theoretic algorithm for constructing, from a group  $G$  of MLF-type, the submodule of  $\bar{k}_+(G)$  which corresponds to the submodule  $(O_k)_+ \subset \bar{k}_+$  of  $\bar{k}_+$  in contradiction to the discussion of (ii). By this fact, one may conclude that

the underlying additive module of the ring of integers of an MLF should be considered to be not group-theoretic .

(Note that if one works with a special case, then a functorial group-theoretic algorithm for constructing the submodule of  $k_+(G)$  which corresponds to the submodule  $(O_k)_+ \subset k_+$  of  $k_+$  may be established. For instance, if  $p(G) \neq 2$ , and  $e(G) = 1$ , then it follows immediately from Lemma 1.2(vi) and Lemma 1.3(iii) that the submodule  $I(G) \subset k_+(G)$  of  $k_+(G)$  in fact corresponds to the submodule  $(O_k)_+ \subset k_+$  of  $k_+$ .)

Remark 4.3.2 .

(i) Let us observe that the field  $\bar{k}$ , as well as the underlying additive module  $\bar{k}_+$  of  $\bar{k}$ , admits the following two natural topologies:

- (a) the  $p_k$ -adic topology
- (b) the natural topology, i.e., the topology determined by the  $p_k$ -adic topologies of the various subfields of  $\bar{k}$  finite over  $k$

Let us also observe that these two topologies do not coincide. One verification of this fact is given as follows: Let us consider the map

$$\text{Tr}: \bar{k}_+ \rightarrow k_+$$



defined by mapping  $a \in \bar{k}_+^*$  to  $d_k^{-1} \text{Tr}_{K=k}(a) \in k_+^*$  if  $a \in K_+^*$  for a finite extension  $K$  of  $k$  contained in  $\bar{k}$  where we write  $\text{Tr}_{K=k}$  for the trace map with respect to the finite extension  $K=k$ . Then one verifies immediately that this map  $\text{Tr}$  is (well-defined and) a  $k$ -linear homomorphism. Moreover, since (one verifies easily that) the restriction of the map  $\text{Tr}$  to each finite extension of  $k$  contained in  $\bar{k}$  is continuous with respect to the respective  $p_k$ -adic topologies on the finite extension and on  $k_+^*$ , the map  $\text{Tr}$  is continuous with respect to the topology (b) on  $\bar{k}_+^*$  and the  $p_k$ -adic topology on  $k_+^*$ .

Next, let us observe that it follows immediately from the well-known surjectivity of the trace map with respect to a finite extension of finite fields that,

for each positive integer  $n$ , there exists an element  $a_n \in (O_{K_n})_+^*$  of  $(O_{K_n})_+^*$  (where we write  $K_n \subset \bar{k}$  for the (uniquely determined) unramified finite extension of  $k$  such that  $d_{K_n} = p^n \cdot d_k$ ) such that  $\text{Tr}_{K_n=k}(a_n) \in O_k \setminus (O_k)_+^*$ .

Then it is immediate that the sequence  $(p^n \cdot a_n)_{n \geq 1}$  converges to  $0 \in \bar{k}_+^*$  with respect to the topology (a) on  $\bar{k}_+^*$ . On the other hand, since  $\text{Tr}(p^n \cdot a_n) = p^{-n} \cdot d_k^{-1} \text{Tr}_{K_n=k}(p^n \cdot a_n) = d_k^{-1} \text{Tr}_{K_n=k}(a_n) \in d_k^{-1} O_k$ ; the sequence  $(\text{Tr}(p^n \cdot a_n))_{n \geq 1}$  does not converge to  $\text{Tr}(0) = 0 \in k_+^*$  with respect to the  $p_k$ -adic topology on  $k_+^*$ . In particular, the map  $\text{Tr}$  is not continuous with respect to the topology (a) on  $\bar{k}_+^*$  and the  $p_k$ -adic topology on  $k_+^*$ . Thus, we conclude that the topology (a) on  $\bar{k}_+^*$  does not coincide with the topology (b) on  $\bar{k}_+^*$ .

(ii) It follows from Summary 3.15 that one has a functorial group-theoretic algorithm for constructing, from a group  $G$  of MLF-type, the topological module  $k_+(G)$ , i.e., the module  $k_+(G)$  equipped with the  $p(G)$ -adic topology. Thus, it follows from the construction of  $\bar{k}_+(G)$  that one has a functorial group-theoretic algorithm for constructing, from a group  $G$  of MLF-type, the module  $\bar{k}_+(G)$  equipped with the natural topology determined by the  $p(G)$ -adic topologies of the various submodules  $k_+(H) = \bar{k}_+(G)^H$  of  $k_+(G)$  where  $H$  ranges over the open subgroups  $H \leq G$  of  $G$ . On the other hand, as already discussed in (i), the  $p_k$ -adic topology on  $\bar{k}_+^*$  does not coincide with the natural topology determined by the  $p_k$ -adic topologies of the various submodules  $k_+ = (\bar{k}_+)^{G_K}$  of  $\bar{k}_+^*$  where  $K$  ranges over the finite extensions of  $k$  contained in  $\bar{k}$ .

(G) the topology on  $\bar{k}_+(G)$  which corresponds to the topology (a) of (i) cf. Remark 4.3.1(i)

(G) the topology on  $\bar{k}_+(G)$  which corresponds to the topology (b) of (i) cf. Summary 3.15, Definition 4.1(i)

(iii) The consideration of the continuity of the map  $\text{Tr}$  of (i) is in fact important in the study of  $p$ -adic Hodge theory. The discussion of (i) asserts that the restriction of the map  $\text{Tr}$  to the (uniquely determined) unramified  $Z_{p_k}$ -extension of  $k$  is not continuous with respect to the respective  $p_k$ -adic topologies. On the other hand, in [12], it was proved that the restriction of the map  $\text{Tr}$  to an arbitrary totally ramified  $Z_{p_k}$ -extension of  $k$  is continuous with respect to the respective  $p_k$ -adic topologies (cf. the discussion preceding [12, Proposition 7]). Moreover, this discontinuity is one of the crucial ingredients of the computation of an important continuous group cohomology  $H^i(G_k; \bar{k}_+(j))$  in the  $p$ -adic Hodge theory established in [12].

Remark 4.3.3 . By Theorem 2.2, Summary 3.15, and Summary 4.3, one may conclude that a group of MLF-type may not be directly related to a sort of ring structure but may be related to one of the underlying additive and underlying multiplicative structures of a ring structure. Put another way, from the point of view of the terminology of [7], a group of MLF-type may/should be regarded as a mono-analytic object (cf. [7, §2.7, (vii)]) (i.e., as opposed to an arithmetic holomorphic object cf. [7, §2.7, (vii)]). Moreover, the passage

from the Galois group  $G_k = \text{Aut}(\bar{k})$  which is, by definition, related to the ring structure of the field  $\bar{k}$

to an abstract group isomorphic to  $G_k$  (i.e., as opposed to a situation in which one regards the group as a group related to the ring structure under consideration as in the case of  $G_k$ )

may be regarded as an analogue of the passage

from the field  $\bar{k}$  equipped with the natural action of  $G_k$

to an abstract monoid isomorphic to either the underlying additive module  $\bar{k}_+$  or the underlying multiplicative monoid  $\bar{k}$  (i.e., as opposed to a situation in which one regards the monoid as a monoid arising from the field  $\bar{k}$ ) equipped with an action of the abstract group isomorphic to  $G_k$ .

### 5. MLF-pairs

One important aspect of mono-anabelian geometry is the technique of mono-anabelian transport (cf. [7, §2]). In order to explain mono-anabelian transport, in the present section, we introduce the notion of an MLF-pair (cf. Definition 5.3 below).

Definition 5.1 .

(i) We shall refer to a collection of data

$$G \curvearrowright M$$

consisting of a monoid  $M$ , a group  $G$ , and an action of  $G$  on  $M$  as a group-monoid-pair.

(ii) Let  $G \curvearrowright M$ ,  $G \curvearrowright M$  be group-monoid-pairs. Then we shall refer to a pair  $= (\phi_G; \phi_M)$  consisting of isomorphisms  $\phi_G: G \xrightarrow{\sim} G$ ,  $\phi_M: M \xrightarrow{\sim} M$  compatible with the respective actions of  $G$ ,  $G$  on  $M$ ,  $M$  as an isomorphism from  $G \curvearrowright M$  to  $G \curvearrowright M$ .

Definition 5.2 . Let  $k$  be an MLF and  $\bar{k}$  an algebraic closure of  $k$ . Then we shall refer to the group-monoid-pair

$$G_k \curvearrowright \bar{k} \quad (\text{respectively, } G_k \curvearrowright O_k^B; \quad G_k \curvearrowright O_k)$$

consisting of the monoid  $\bar{k}$  (respectively,  $O_k^B; O_k$ ), the group  $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=k)$ , and the natural action of  $G_k$  on  $\bar{k}$  (respectively,  $O_k^B; O_k$ ) as the model MLF-pair (respectively, model MLF<sup>B</sup>-pair; model MLF-pair) (associated to  $\bar{k}=k$ ).

Definition 5.3 . We shall refer to an isomorph, as a group-monoid-pair (i.e., in the sense of Definition 5.1 (ii)), of a model MLF -pair (respectively, model MLF<sup>B</sup>-pair; model MLF -pair) as an MLF -pair (respectively, MLF<sup>B</sup>-pair; MLF -pair).

Remark 5.3.1 . The notion of an MLF -pair (respectively, MLF<sup>B</sup>-pair; MLF -pair) in the sense of Definition 5.3 is the same as the notion of an MLF-Galois<sup>TM</sup>-pair of mono-analytic type (respectively, MLF-Galois TCG-pair of mono-analytic type) in the sense of [6, Definition 3.1 (ii)] (cf. also Remark 5.7.2 below).

In the remainder of the present section, let  $f : B \rightarrow g$  and  $G \rightarrow M$  an MLF -pair.

Definition 5.4 . We shall refer to the group  $G$  (respectively, monoid  $M$ ) as the étale-like portion (cf. [7, §2.7 (iii)]) (respectively, Frobenius-like portion (cf. [7, §2.7 (ii)])) of the MLF -pair  $G \rightarrow M$ .

Remark 5.4.1 . It follows from the various definitions involved that the étale-like portion of an MLF -pair is a group of MLF-type.

Definition 5.5 . We shall write

$$\text{Ind} \stackrel{\text{def}}{=} \begin{cases} \sum_{\mathbb{Z}} f^{-1}g & \text{if } = \\ \prod_{\mathbb{Z}} f^{-1}g & \text{if } = B \\ \prod_{\mathbb{Z}} & \text{if } = : \end{cases}$$

Let us regard  $\text{Ind}$  as a subgroup of  $\mathbb{Z}$  (in an evident way).

Remark 5.5.1 . Let us observe that, for each element of  $\text{Ind}$ , the identity automorphism of  $G$  and the automorphism of  $M$  given by multiplication by the element of  $\text{Ind}$  (i.e., the automorphism given by  $x \mapsto x$  for the  $x \in \mathbb{Z} \subset \text{Ind} \subset \mathbb{Z}$ ) form an automorphism of the MLF -pair  $G \rightarrow M$ . In particular, we have an injective homomorphism

$$\text{Ind} \hookrightarrow \text{Aut}_G(M) \subset \text{Aut}(G \rightarrow M) :$$

Lemma 5.6 . Suppose that  $=$ . Then the following hold:

- (i) Let  $H \subset G$  be an open subgroup of  $G$ . Then there exist infinitely many prime numbers  $l$  such that the pro- $l$  completion of the submodule  $M^H \subset M$  of  $H$ -invariants of  $M$  is, as an abstract pro- $l$  group, isomorphic to  $(\mathbb{Z}_l)_+$ . Moreover, for such an  $l$ , the submodule

$$J(H) \subset M^H$$

of  $M^H$  obtained by forming the kernel of the natural homomorphism from  $M^H$  to the pro- $l$  completion of  $M^H$  does not depend on the choice of  $l$ .

- (ii) Write

$$M(\ ) \stackrel{\text{def}}{=} \varprojlim_{H \subset G} J(H) \subset M$$

(cf. (i)) where the injective limit is taken over the open subgroups  $H \subset G$  of  $G$ . Then the submodule  $M(\ ) \subset M$  of  $M$  is  $G$ -stable.

Proof. Let us observe that it follows from the various definitions involved that, for each open subgroup  $H \leq G$  of  $G$ , the module  $M^H$  is, as an abstract module, isomorphic to  $k$  for some MLF. Thus, these assertions follow immediately from Lemma 1.2(i).

Proposition 5.7 . The following hold:

- (i) Suppose that  $\mathcal{O} = B$ . Then the group-monoid-pair  $G \curvearrowright M^{gp}$  obtained by replacing  $M$  in  $G \curvearrowright M$  by  $M^{gp}$  is an MLF -pair.
- (ii) Suppose that  $\mathcal{O} = \bar{k}$ . Then the group-monoid-pair  $G \curvearrowright M(\bar{k})$  obtained by replacing  $M$  in  $G \curvearrowright M$  by  $M(\bar{k})$  of Lemma 5.6(ii) is an MLF -pair.

Proof. Assertion (i) follows from the (easily verified) fact that the natural inclusion  $\mathcal{O}_k^B \hookrightarrow \bar{k}$  determines an isomorphism  $(\mathcal{O}_k^B)^{gp} \cong \bar{k}$ . Assertion (ii) follows immediately in light of Lemma 1.2(i) from the definition of  $M(\bar{k})$ .

Remark 5.7.1 . The content of Proposition 5.7 may be summarized as follows:

$$\text{an MLF}^B\text{-pair } (G \curvearrowright M) \Rightarrow \text{an MLF-pair } (G \curvearrowright M^{gp}) \Rightarrow \text{an MLF-pair } (G \curvearrowright M(\bar{k})).$$

Remark 5.7.2 .

- (i) Suppose that  $\mathcal{O} = \bar{k}$ . Then it follows from the various definitions involved that, for each open subgroup  $H \leq G$  of  $G$ , the module  $M^H$  corresponds to the module  $\mathcal{O}$  for some MLF. Thus, by considering the submodules of finite index of  $M^H$ , one obtains a topology on  $M^H$  which corresponds to the  $p$ -adic topology on  $\mathcal{O}$ . In particular, one may regard  $M$  as an object of the category  $\text{TCG}$  of [6, Definition 3.1 (i)].
- (ii) Suppose that  $\mathcal{O} = k$  (respectively,  $\mathcal{O} = B$ ). Then it follows from the various definitions involved that, for each open subgroup  $H \leq G$  of  $G$ , the submonoid  $M^H$  of  $H$ -invariants of  $M$  contains the submodule  $M^{gp}(k)^H$  of  $H$ -invariants of  $M^{gp}(k)$  (cf. Proposition 5.7(i), (ii)); moreover, the pair of monoids  $M^{gp}(k)^H \leq M^H$  corresponds to the pair of monoids  $\mathcal{O} \leq k$  (respectively,  $\mathcal{O} \leq B$ ) for some MLF. Thus, by considering the topology on  $M^H$  induced by the topology of  $M^{gp}(k)^H$  discussed in (i) (cf. Proposition 5.7(i), (ii)), one obtains a topology on  $M^H$  which corresponds to the  $p$ -adic topology on  $k$  (respectively,  $\mathcal{O}^B$ ). In particular, one may regard  $M$  as an object of the category  $\text{TLG}$  (respectively,  $\text{TM}$ ) of [6, Definition 3.1 (i)].

Definition 5.8 . We shall write

$$M(G) \stackrel{\text{def}}{=} \begin{cases} \mathcal{O} \otimes \bar{k}(G) & \text{if } \mathcal{O} = \bar{k} \\ \mathcal{O}^B(G) & \text{if } \mathcal{O} = B \\ \bar{\mathcal{O}}(G) & \text{if } \mathcal{O} = k \end{cases}$$

(cf. Definition 4.1 (i), Remark 5.4.1). Thus, it follows from Proposition 4.2(i) that the group-monoid-pair

$$G \curvearrowright M(G)$$

is an MLF -pair.

Definition 5.9 .

(i) We shall refer to the group-monoid-pair

$$G_y (G)$$

(cf. Definition 4.1 (iii), Remark 5.4.1), i.e., obtained by considering the cyclotome associated to  $G$ , as the étale-like cyclotome associated to  $G_y M$ .

(ii) We shall write

$$(M) \stackrel{\text{def}}{=} \varinjlim_n M[n] = \varinjlim_n M^{\text{gp}[n]}$$

where the projective limits are taken over the positive integers  $n$  and refer to the group-monoid-pair

$$G_y (M)$$

obtained by considering the action of  $G$  on  $(M)$  induced by the action of  $G$  on  $M$  as the Frobenius-like cyclotome associated to  $G_y M$ . We regard the cyclotome  $(M)$  as a pro finite, hence also topological,  $G$ -module (by the easily verified finiteness of  $M[n]$  for each  $n$ ).

Remark 5.9.1 . One verifies easily from the various definitions involved that we have a natural identification of the étale-like cyclotome  $G_y (G)$  with the Frobenius-like cyclotome  $G_y (M(G))$  associated to the MLF -pair  $G_y M(G)$  constructed from  $G$  in Definition 5.8.

Proposition 5.10 . Let  $k$  be an MLF and  $\bar{k}$  an algebraic closure of  $k$ . Suppose that the MLF -pair  $G_y M$  is the model MLF -pair associated to  $\bar{k}=k$  (which thus implies that  $G = G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}=k)$  and  $M = \bar{k}$ ). Then the following hold:

(i) The isomorphism  $(\bar{k})! (G)$  of topological  $G$ -modules of Proposition 4.2(iv) induces an isomorphism of the étale-like cyclotome

$$G = G_k y (G)$$

with the group-monoid-pair

$$G_k y (\bar{k})$$

obtained by considering the cyclotome associated to  $\bar{k}$ .

(ii) The natural inclusion  $(\bar{k})! M$  induces an isomorphism of the Frobenius-like cyclotome

$$G = G_k y (M)$$

with the group-monoid-pair

$$G_k y (\bar{k})$$

obtained by considering the cyclotome associated to  $\bar{k}$ .

Proof. These assertions are immediate.

### 6. Cyclotomic Synchronization for MLF-pairs

Let us recall that a cyclotome refers to an isomorph of the object  $\mathcal{Z}(1)$ . Various (a priori independent) cyclotomes often appear in studies of arithmetic geometry. For instance, if  $k$  is an algebraically closed field of characteristic zero, then each of

the cyclotome  $(\mathcal{Z})$  associated to  $k$ ,

the étale fundamental group

$$\pi_1^{\text{ét}}(\text{Spec}(k((t))))$$

of the spectrum of the field  $k((t))$  of fractions of the ring of formal power series with coefficients in  $k$ , and

the dual

$$(\mathcal{Z}) \stackrel{\text{def}}{=} \text{Hom}_{\mathbb{Z}}(H_{\text{ét}}^2(C; \mathbb{Z}); \mathbb{Z})$$

over  $\mathbb{Z}$  of the second étale cohomology  $H_{\text{ét}}^2(C; \mathbb{Z})$  of a projective smooth curve  $C$  over  $k$

gives an example of a cyclotome. In our case, we constructed, from a single MLF-pair  $G \rightarrow M$  (where  $2f \in B; g$ ), (a priori independent) two cyclotomes, i.e.,

the étale-like cyclotome  $(\mathcal{Z}(G))$  and

the Frobenius-like cyclotome  $(\mathcal{Z}(M))$

(cf. Definition 5.9, Proposition 5.10).

If one works with certain arithmetic holomorphic structures (cf. [7, §2.7, (vii)]) related to cyclotomes under consideration, then one may obtain a phenomenon of cyclotomic synchronization, i.e., synchronization of cyclotomes. For instance, in the case of the above examples  $(\mathcal{Z}(G))$  and  $(\mathcal{Z}(M))$ , the homomorphism  $\text{Pic}(C) \rightarrow H^2(C; \mathbb{Z})$  obtained by considering the first Chern classes yields an isomorphism

$$\text{Pic}(C) \cong \text{Pic}^0(C) \cong \mathbb{Z} \oplus \text{Hom}_{\mathbb{Z}}(\mathcal{Z}(C); \mathbb{Z});$$

thus, an invertible sheaf on  $C$  of degree one determines a cyclotomic synchronization (i.e., an isomorphism between cyclotomes)  $(\mathcal{Z}(C)) \cong (\mathcal{Z}(M))$  by means of which one usually (i.e., in the usual point of view of arithmetic geometry) identifies  $(\mathcal{Z}(C))$  with  $(\mathcal{Z}(M))$ . In our case, if our MLF-pair  $G \rightarrow M$  is a model MLF-pair (i.e., an MLF-pair directly related to the ring structure of an MLF), then, by means of the arithmetic holomorphic structures related to the objects involved (more concretely, essentially, by means of local class field theory), one may construct a natural isomorphism between the Frobenius-like cyclotome  $(\mathcal{Z}(M))$  and the étale-like cyclotome  $(\mathcal{Z}(G))$  (cf. Definition 6.2 below).

On the other hand, observe that such a phenomenon of cyclotomic synchronization usually depends at least in an a priori sense, on a certain arithmetic holomorphic structure related to cyclotomes under consideration. Thus, if one works with only mono-analytic structures (cf. [7, §2.7, (vii)]), then one cannot apply the arithmetic holomorphic construction of the cyclotomic synchronization. In the present section, we establish a cyclotomic synchronization that one may apply to an arbitrary (i.e., a not necessarily model) MLF-pair (cf. Definition 6.6, Proposition 6.7 below).

In the present section, let  $2f \in B; g$  and  $G \rightarrow M$  an MLF-pair.

Proposition 6.1 . The set

$$\text{Isom}_G ( M ) ; ( G )$$

of  $G$ -equivariant isomorphisms  $( M ) ! ( G )$  (is nonempty and) forms a  $\mathcal{Z}$ -torsor, i.e., relative to the natural action of  $\mathcal{Z} = \text{Aut}( ( G ) )$  on  $( G )$ .

Proof. This assertion follows immediately from Proposition 5.10(i), (ii).

De nition 6.2 . Suppose that the MLF -pair  $G y M$  is a model MLF -pair. Then, by considering the composite of the isomorphisms of Proposition 5.10(i), (ii) (i.e, the isomorphisms  $( G ) ! ( \bar{k} )$ ,  $( M ) ! ( \bar{k} )$ ), we obtain a  $G$ -equivariant isomorphism

$$( M ) ! ( G ) :$$

We shall refer to this composite  $( M ) ! ( G )$  (that is an element of the  $\mathcal{Z}$ -torsor discussed in Proposition 6.1) as the holomorphic cyclotomic synchronization isomorphism associated to the model MLF -pair  $G y M$ .

Remark 6.2.1 . One veri es easily that the construction of the holomorphic cyclotomic synchronization isomorphism of De nition 6.2 depends in an essential way, on the ring structure (i.e., the arithmetic holomorphic structure cf. [7, §2.7, (vii)]) of a given model MLF -pair. In particular, one cannot apply this construction of the holomorphic cyclotomic synchronization isomorphism of De nition 6.2 to an abstract (i.e., a not necessarily model) MLF -pair.

Lemma 6.3 . Suppose that  $2 f ; Bg$ . Write  $M ( ) \stackrel{\text{def}}{=} M^{gp} ( )$  (cf. Proposition 5.7(i), (ii)). (Thus, one veri es easily that  $M ( ) = M = M^{gp}$ .) Write, moreover,  $V(G y M) \stackrel{\text{def}}{=} M^{l(G)} = M ( )^{l(G)}$  (cf. De nition 3.5(iii), Remark 5.4.1). (Thus, one also veri es easily that there exists a natural identi cation  $V(G y M)^{gp} = (M^{gp})^{l(G)} = M ( )^{l(G)}$ .) Then the following hold:

(i) The monoid  $V(G y M)$  is, as an abstract monoid, isomorphic to

$$\begin{cases} \mathbb{Z}_+ & \text{if } = \\ \mathbb{N} & \text{if } = B: \end{cases}$$

Moreover, the action of  $G$  on  $V(G y M)$  induced by the action of  $G$  on  $M$  is trivial .

(ii) For each positive integern, the natural homomorphism

$$H^2(G; M [n]) ! H^2(G; M^{gp})$$

determines an isomorphism

$$H^2(G; M [n]) ! H^2(G; M^{gp})[n]:$$

(iii) The natural homomorphism

$$H^2 G=l(G); (M^{gp})^{l(G)} ! H^2(G; M^{gp})$$

is an isomorphism

(iv) The natural homomorphism

$$H^2(G; (M^{gp})^{I(G)}) \rightarrow H^2(G; V(G, M)^{gp})$$

is an isomorphism.

(v) The homomorphism

$$H^1(G; V(G, M)^{gp}) \xrightarrow{pf} V(G, M)^{gp} \rightarrow H^2(G; V(G, M)^{gp})$$

determined by the exact sequence of  $\mathbb{Z}[G]$ -modules

$$0 \rightarrow V(G, M)^{gp} \rightarrow V(G, M)^{gp} \xrightarrow{pf} V(G, M)^{gp} \xrightarrow{pf} V(G, M)^{gp} \rightarrow 0$$

is an isomorphism.

(vi) The homomorphism

$$H^1(G; V(G, M)^{gp}) \xrightarrow{pf} V(G, M)^{gp} \rightarrow V(G, M)^{gp} \xrightarrow{pf} V(G, M)^{gp}$$

obtained by mapping

$$\begin{aligned} 2 \text{ Hom}(G; V(G, M)^{gp}) \xrightarrow{pf} V(G, M)^{gp} \\ = H^1(G; V(G, M)^{gp}) \xrightarrow{pf} V(G, M)^{gp} \end{aligned}$$

(cf. (i)) to

$$\text{Frob}(G) \xrightarrow{2} V(G, M)^{gp} \xrightarrow{pf} V(G, M)^{gp}$$

(cf. Definition 3.8, Remark 5.4.1) is an isomorphism.

(vii) The various isomorphisms of (iii), (iv), (v), (vi) determine an isomorphism

$$H^2(G; M^{gp}) \rightarrow V(G, M)^{gp} \xrightarrow{pf} V(G, M)^{gp}$$

which thus determines an isomorphism

$$H^2(G; (M)) \rightarrow V(G, M)^{gp}$$

(cf. (i), (ii)).

**Proof.** These assertions follow in light of Proposition 3.6 and Proposition 3.9 from Lemma 1.6 (i), (ii), (iii), (iv), (v), (vi), (vii).

**Definition 6.4 .** Suppose that  $2 \text{ f } ; Bg$ . Then we shall write

$$\text{inv}^\wedge(G, M): H^2(G; (M)) \rightarrow V(G, M)^{gp}$$

for the second displayed isomorphism of Lemma 6.3 (vii).

**Lemma 6.5 .** Suppose that  $2 \text{ f } ; Bg$ . Recall the  $\mathbb{Z}$ -torsor

$$\text{Isom}_G((M); (G))$$



discussed in Proposition 6.1. Then the subset of  $\text{Isom}_G((M); (G))$  consisting of isomorphisms  $\in \text{Isom}_G((M); (G))$  which satisfy the following condition (is nonempty and forms an  $\text{Ind}$ -torsor: The composite

$$V(G y M)^{\text{gp}} \xrightarrow{\text{inv}^\wedge(G y M)} H^2(G; (M)) \xrightarrow{H^2(\text{id}_G; \cdot)} H^2(G; (G)) \xrightarrow{\text{inv}^\wedge(G y M (G))} V(G y M (G))^{\text{gp}}$$

(cf. Remark 5.9.1) where we write  $H^2(\text{id}_G; \cdot)$  for the isomorphism induced by maps the submonoid

$$V(G y M) \rightarrow V(G y M)^{\text{gp}}$$

(cf. Lemma 6.3(i)) to the submonoid

$$V(G y M (G)) \rightarrow V(G y M (G))^{\text{gp}}$$

**Proof.** Let us first observe that one verifies immediately from Proposition 5.10(i), (ii) that the subset discussed in Lemma 6.5 is nonempty. Next, let us observe that, by Lemma 6.3(i), there exists a natural isomorphism of  $\text{Aut}(V(G y M))$  with  $\text{Ind}$ . Thus, it follows immediately from the definition of the displayed composite in the statement of Lemma 6.5 that the subset discussed in Lemma 6.5 forms an  $\text{Ind}$ -torsor, as desired.

**Definition 6.6.** If  $\mathcal{C} = \mathcal{C}$ , then we shall write

$$\text{syn}(G y M) \stackrel{\text{def}}{=} \text{Isom}_G((M); (G)) :$$

If  $\mathcal{C} = \mathcal{C}; Bg$ , then we shall write

$$\text{syn}(G y M) = \text{Isom}_G((M); (G))$$

for the  $\text{Ind}$ -torsor discussed in Lemma 6.5. (So, for each  $\mathcal{C} = \mathcal{C}; Bg$ , the set  $\text{syn}(G y M)$  forms an  $\text{Ind}$ -torsor cf. also Proposition 6.1). We shall refer to the  $\text{Ind}$ -torsor  $\text{syn}(G y M)$  as the cyclotomic synchronization poly-isomorphism associated to  $G y M$ .

**Proposition 6.7.** Suppose that the MLF-pair  $G y M$  is a model MLF-pair. Then the  $\text{Ind}$ -torsor  $\text{syn}(G y M)$  coincides with the  $\text{Ind}$ -orbit of the holomorphic cyclotomic synchronization isomorphism associated to  $G y M$ .

**Proof.** This assertion follows immediately from the definition of  $\text{syn}(G y M)$ .

**Remark 6.7.1.**

- (i) Let us observe that one verifies easily from the definition of the injective homomorphism of Remark 5.5.1 that the composite

$$\text{Ind} \rightarrow \text{Aut}(G y M) \rightarrow \text{Aut}(G y (M)) \rightarrow \text{Aut}((M)) = \mathcal{C}$$

coincides with the natural inclusion  $\text{Ind} \rightarrow \mathcal{C}$ , but the image of the composite

$$\text{Ind} \rightarrow \text{Aut}(G y M) \rightarrow \text{Aut}(G) \rightarrow \text{Aut}((G)) = \mathcal{C}$$

consists of  $\mathcal{C}$ .

(ii) Let

$$s : G^y y M^y \rightarrow s (G^y y M^y) \text{ Isom}_{G^y} ( M^y); ( G^y)$$

be a functorial assignment which assigns each MLF-pair  $G^y y M^y$  (for a fixed  $2 f ; B; g$ ) to a subset of the  $\mathbb{Z}$ -torsor  $\text{Isom}_{G^y}(( M^y); ( G^y))$ . Then, by the observation of (i), one may verify easily that

the subset  $s (G^y y M^y) \text{ Isom}_{G^y}(( M^y); ( G^y))$  is stable by the action of  $\text{Ind } \mathbb{Z}$ .

(iii) Let us recall that the cyclotomic synchronization poly-isomorphism  $\text{syn} (G y M)$  (which forms by allowing  $G y M$  to vary a functorial assignment as discussed in (ii))

( ) contains the holomorphic cyclotomic synchronization isomorphism (i.e., if one considers the cyclotomic synchronization poly-isomorphism of a model MLF -pair) (cf. Proposition 6.7).

By the observation of (ii), together with Proposition 6.7, one may conclude that the cyclotomic synchronization poly-isomorphism  $\text{syn} (G y M)$  (which forms by allowing  $G y M$  to vary a functorial assignment as discussed in (ii)) is minimal among functorial assignments as discussed in (ii) which satisfy the condition ( ).

Put another way, roughly speaking, the cyclotomic synchronization poly-isomorphism of Definition 6.6 may be considered to be best among functorial assignments as discussed in (ii).

### 7. Mono-anabelian Transport for MLF-pairs

One important aspect of the technique of mono-anabelian transport is the notion of a Kummer poly-isomorphism, which relates (monoids constructed, via some functorial algorithms, from) Frobenius-like portions to mono-anabelian étale-like monoids (i.e., monoids constructed, via some mono-anabelian reconstruction algorithms, from étale-like portions). In the present section, we establish Kummer poly-isomorphisms (cf. Definition 7.4 below) and discuss mono-anabelian transport for MLF-pairs (cf. Remark 7.6.1 below).

In the present section, let  $2 f ; B; g$  and  $G y M$  an MLF -pair.

Lemma 7.1 . The following hold:

(i) For each open subgroup  $H \subset G$  of  $G$ , the homomorphism

$$(M^H) \rightarrow (M^{gp})^H \rightarrow H^1 H; ( M)$$

determined by the exact sequences of  $\mathfrak{bf}$ -modules

$$1 \rightarrow M^{gp}[n] \rightarrow M^{gp} \rightarrow 1$$

where  $n$  ranges over the positive integers is injective.

(ii) The homomorphism

$$(M) \rightarrow M^{gp} \rightarrow H^1 G; ( M)$$

obtained by forming the injective limit of the homomorphisms of (i) for the open subgroups  $H \subset G$  is injective.

Proof. First, we verify assertion (i). Let us first observe that, by definition, the kernel of the homomorphism discussed in assertion (i) coincides with the submodule  $\alpha(M^{gp})^H$  consisting of divisible elements of  $(M^{gp})^H$ . Next, let us observe that the module  $(M^{gp})^H$  is, as an abstract module, isomorphic to  $k$  (respectively,  $k; O$ ) for some MLF if  $\alpha =$  (respectively,  $\alpha = B; \alpha =$ ). Thus, assertion (i) follows from Lemma 1.2(i). This completes the proof of assertion (i). Assertion (ii) follows from assertion (i).

Definition 7.2 . We shall write

$${}_1 \text{Kmm}(G y M): M \rightarrow {}_1 H^1 G; (M)$$

for the injection (obtained by the injection) discussed in Lemma 7.1(ii).

Lemma 7.3 . The  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms

$${}_1 H^1 G; (M) \rightarrow {}_1 H^1 G; (G)$$

induced by the  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms

$$\text{syn}(G y M): (M) \rightarrow (G)$$

restricts relative to the injections  ${}_1 \text{Kmm}(G y M)$ ,  ${}_1 \text{Kmm}(G y M (G))$  of Definition 7.2 (cf. also Remark 5.9.1) to an  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms

$$M \rightarrow M (G):$$

Proof. This assertion follows immediately from Proposition 6.7.

Definition 7.4 . We shall write

$$(G y M): (G y M) \rightarrow G y M (G)$$

for the  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms discussed in Lemma 7.3 and refer to  $(G y M)$  as the Kummer poly-isomorphism associated to  $G y M$ .

Some of the various assertions discussed in Sections 5, 6, and 7 may be summarized as follows.

Summary 7.5 . One may construct, from an MLF -pair  $G y M$ ,

the étale-like MLF -pair  $G y M (G)$ ,

the étale-like cyclotome  $G y (G)$ ,

the Frobenius-like cyclotome  $G y (M)$ ,

the cyclotomic synchronization poly-isomorphisms  $\text{syn}(G y M): (M) \rightarrow (G)$  (i.e., a certain poly-isomorphism which forms an  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms  $(M) \rightarrow (G)$ ), and

the Kummer poly-isomorphism  $(G y M): M \rightarrow M (G)$  (i.e., a certain poly-isomorphism which forms an  $\text{Ind}$ -orbit of  $G$ -equivariant isomorphisms  $M \rightarrow M (G)$ ).

**Theorem 7.6.** — Let  $(G, B, \times)$ ;  $G \curvearrowright M, G \cdot \curvearrowright M$ . MLF -pairs. Thus, we have a natural map

$$F: \text{Isom}(G \curvearrowright M, G \cdot \curvearrowright M) \rightarrow \text{Isom}(G, G).$$

Moreover, by considering the action of  $\text{Ind}$  on  $M$ . (cf. Remark 5.5.1), we have a natural action of  $\text{Ind}$  on the set  $\text{Isom}(G \curvearrowright M, G \cdot \curvearrowright M)$  over  $\text{Isom}(G, G)$  relative to the map  $F$ . Then the following hold:

- (i) The map  $F$  is surjective.
- (ii) Every fiber of the map  $F$  forms an  $\text{Ind}$  -torsor.
- (iii) If  $\times = \cdot$ , then every fiber of the map  $F$  is of cardinality two.
- (iv) If  $\times = B$ , then the map  $F$  is bijective.

*Proof.* — First, we verify assertion (i). Let  $\gamma : G \rightarrow G$ . be an isomorphism (i.e., an element of the codomain of  $F$ ). Then one verifies easily from the *functoriality* of the mono-anabelian reconstruction algorithm of Definition 4.1(i) that  $\gamma$  induces an isomorphism between the “étale-like” MLF -pairs

$$G \curvearrowright M (G) \xrightarrow{\gamma, M(\gamma)} G \cdot \curvearrowright M (G).$$

Thus, by considering respective elements  $\gamma, \gamma \cdot$  of the Kummer poly-isomorphisms  $(G \curvearrowright M), (G \cdot \curvearrowright M)$ , we obtain isomorphisms

$$(G \curvearrowright M) \xrightarrow{\gamma, M(\gamma)} G \curvearrowright M (G) \xrightarrow{\gamma \cdot, M(\gamma \cdot)} G \cdot \curvearrowright M (G) \xrightarrow{\gamma \cdot, M(\gamma \cdot)} (G \cdot \curvearrowright M)$$

such that the image of the composite  $(G \curvearrowright M) \rightarrow (G \cdot \curvearrowright M)$  of these isomorphisms via  $F$  coincides with the original isomorphism  $\gamma$ . This completes the proof of assertion (i). Next, we verify assertion (ii). Let us first observe that, to verify assertion (ii), by considering the difference of two elements of the domain of  $F$  whose images via  $F$  coincide, it suffices to verify the following assertion:

Let  $(\gamma, M)$  be an automorphism of the MLF -pair  $G \curvearrowright M$ . Suppose that  $\gamma = \text{id}_G$ . Then it holds that  $M \in \text{Ind}(\text{Aut}_G(M))$ .

To this end, let us observe that one verifies easily from the *functoriality* of the Kummer poly-isomorphism of Definition 7.4 that the Kummer poly-isomorphism associated to  $G \curvearrowright M$

$$(G \curvearrowright M): (G \curvearrowright M) \rightarrow G \curvearrowright M (G)$$

is compatible with the automorphism  $(\gamma, M)$ , i.e., the diagram of *poly-isomorphisms*


$$\begin{array}{ccc} (G \curvearrowright M) & \xrightarrow{(\gamma, M)} & G \curvearrowright M (G) \\ (\gamma, M) \downarrow & & \downarrow \gamma, M(\gamma) \\ (G \curvearrowright M) & \xrightarrow{(\gamma, M)} & G \curvearrowright M (G) \end{array}$$

commutes. Next, let us observe that since (we have assumed that)  $\text{id}_G = \text{id}_G$ , the right-hand vertical arrow of this diagram is the (poly-isomorphism consisting of the) *identity automorphism*. Thus, it holds that  $(\text{id}_G, M) = (G \curvearrowright M)^{-1} (G \curvearrowright M)$ , which thus implies that  $\text{Ind}_M = \text{Ind}(\text{Aut}_G(M))$ , as desired. This completes the proof of assertion (ii). Assertions (iii), (iv) follow from assertions (i), (ii), together with the (easily verified) fact that  $\text{Ind} = 2$  and  $\text{Ind}^B = 1$ .

**Remark 7.6.1.** — The content of Theorem 7.6, as well as the proof of Theorem 7.6, gives some examples of the technique of *mono-anabelian transport*.

- (i) In order to explain the technique of *mono-anabelian transport* from the point of view of Theorem 7.6 (i), let us recall the proof of Theorem 7.6 (i), i.e., the *surjectivity* of the map  $F$  of Theorem 7.6 as follows:


Theorem 7.6 (i) asserts that, roughly speaking, for two MLF -pairs, an “étale-like link” between the MLF -pairs (i.e., an isomorphism between the *étale-like portions* of the MLF -pairs) induces a “Frobenius-like link” between the MLF -pairs (i.e., an isomorphism between the *Frobenius-like portions* of the MLF -pairs) compatible with the given “étale-like link”.



Suppose that we are in a situation described by the following diagram to recall the proof of Theorem 7.6 (i):




In order to obtain a “Frobenius-like link” from our “étale-like link”, let us first apply the *mono-anabelian reconstruction algorithm* discussed in Summary 4.3 to each of the étale-like portions to construct a mono-anabelian étale-like monoid (i.e., the “étale-like copy  $M(G)$ ” — cf. Definition 5.8 — of the Frobenius-like portion “ $M$ ”).




Here, let us recall that we do *not require* the “étale-like link” to be *compatible* with any sort of ring structures. On the other hand, observe that the mono-anabelian reconstruction algorithms discussed in Sections 3 and 4 may be applied *without* existence of some “fixed reference model” (as  $\bar{k}/k$  and  $\bar{k}_*/k_*$  for  $G = \text{Gal}(\bar{k}/k)$  and  $G_* = \text{Gal}(\bar{k}_*/k_*)$  in the case of bi-anabelian geometry discussed in Section 2). Put another way, the mono-anabelian reconstruction algorithms discussed in Sections 3 and 4 have the virtue of being *free* of any mention of some “fixed reference model” copy of ring-theoretic objects. In particular, by the mono-anabelian property of our algorithm, we obtain, from the given “étale-like link”, an isomorphism between the mono-anabelian étale-like monoids


compatible with the given “étale-like link”.




Next, in order to relate the Frobenius-like portions to the mono-anabelian étale-like monoids, let us discuss *cyclotomic synchronization poly-isomorphisms*, which induce *Kummer poly-isomorphisms*. Let us recall that, as in the discussion of Section 6, we can construct a *cyclotomic synchronization poly-isomorphism* “syn” (cf. Definition 6.6) (i.e., a poly-isomorphism between the Frobenius-like cyclotome and the étale-like cyclotome).



Thus, by applying a sort of the *Kummer theory* (i.e., by considering the injection “Kmm” — cf. Definition 7.2), we obtain a *Kummer poly-isomorphism* “K” (cf. Definition 7.4) (i.e., a poly-isomorphism between the Frobenius-like portion and the mono-anabelian étale-like monoid) from the above cyclotomic synchronization poly-isomorphism.



Thus, we are in a situation described by the following diagram:




In particular, by considering the composite of the two lower vertical arrows and the middle horizontal arrow of this diagram, we obtain a “Frobenius-like link” compatible with the given “étale-like link”.

Thus, in summary, by means of

- *Kummer-detachment* (cf. [7, §2.7, (vi)]), i.e., the passage, via Kummer poly-isomorphisms, from Frobenius-like structures to corresponding étale-like structures, and
- *étale-transport*, i.e., the passage, via the mono-anabelian property of mono-anabelian reconstruction algorithms, from the “left-hand side” to the “right-hand side”,

one may *transport*, via “étale-like links”, Frobenius-like portions from the “left-hand side” to the “right-hand side”.



(ii) In order to explain the technique of *mono-anabelian transport* from the point of view of Theorem 7.6 (ii), let us recall the proof of Theorem 7.6 (ii) as follows:

Theorem 7.6 (ii) asserts that, roughly speaking, for an isomorphism between MLF -pairs, one may compute the effect on the Frobenius-like portions of the “Frobenius-like link” (cf. (i)) from the point of view of the effect on the étale-like portions of the “étale-like link” (cf. (i)). Suppose that we are in a situation described by the following diagram to recall the proof of Theorem 7.6 (ii):

$$\begin{array}{ccc} \text{étale-like portion} & - & \text{étale-like portion} \\ \downarrow \gamma & & \downarrow \gamma \\ \text{Frobenius-like portion} & - & \text{Frobenius-like portion.} \end{array}$$

In order to compute the effect of the “Frobenius-like link” from the point of view of the effect of the “étale-like link”, let us first apply the *mono-anabelian reconstruction algorithm* discussed in Summary 4.3 to each of the étale-like portions to construct a mono-anabelian étale-like monoid (cf. (i)).

$$\text{étale-like portion} \xrightarrow[\text{reconstruction algorithm}]{\text{mono-anabelian}} \text{mono-anabelian étale-like monoid}$$


Here, let us recall that we do *not require* the “links” to be *compatible* with any sort of ring structures. On the other hand, observe that the mono-anabelian reconstruction algorithms discussed in Sections 3 and 4 may be applied *without* existence of some “fixed reference model” (as  $\bar{k}/k$  and  $\bar{k}_./k_.$  for  $G = \text{Gal}(\bar{k}/k)$  and  $G_ = \text{Gal}(\bar{k}_./k_.)$  in the case of bi-anabelian geometry discussed in Section 2). Put another way, the mono-anabelian reconstruction algorithms discussed in Sections 3 and 4 have the virtue of being *free* of any mention of some “fixed reference model” copy of ring-theoretic objects. In particular, by the mono-anabelian property of our algorithm, we obtain, from the “étale-like link”, an isomorphism between the mono-anabelian étale-like monoids *compatible* with the “étale-like link”.

$$\begin{array}{ccc} \text{étale-like portion} & - & \text{étale-like portion} \\ \downarrow \gamma & \text{mono-anab. property} & \downarrow \gamma \\ \text{mono-anabelian étale-like monoid} & - & \text{mono-anabelian étale-like monoid} \end{array}$$

Next, in order to relate the Frobenius-like portions to the mono-anabelian étale-like monoids, let us discuss *cyclotomic synchronization poly-isomorphisms*, which induce *Kummer poly-isomorphisms*. As in the discussion of (i), by applying a sort of the *Kummer theory*, we obtain a *Kummer poly-isomorphism* “ ” from the cyclotomic synchronization poly-isomorphism “syn ”.

$$\begin{array}{ccc} & \text{cyclotomic} & \\ & \text{synchronization} & \\ \text{Frobenius-like cyclotome} & - & \text{étale-like cyclotome} \\ & \text{Kummer} & \\ \text{Kummer theory} & \text{poly-isomorphism} & \\ \text{Frobenius-like portion} & - & \text{mono-anabelian étale-like monoid} \end{array}$$

Thus, we are in a situation described by the following diagram:




Here, let us observe that, *in our case*, by the constructions of the various objects under consideration, the lower square (of *poly-isomorphisms*) of this diagram *commutes*. In particular, by this *commutativity*, we can compute the effect on the Frobenius-like portions of the “Frobenius-like link” by considering the composite of the Kummer poly-isomorphism, a(n) (poly-)isomorphism that arises from the mono-anabelian property of the mono-anabelian reconstruction algorithm, and the inverse of the Kummer poly-isomorphism.

Thus, in summary, by means of

- *Kummer-detachment*, i.e., the passage, via Kummer poly-isomorphisms, from Frobenius-like structures to corresponding étale-like structures, and
- *étale-transport*, i.e., the passage, via the mono-anabelian property of mono-anabelian reconstruction algorithms, from the “left-hand side” to the “right-hand side”,

one may *compute* the effect of the “Frobenius-like link” from the point of view of the effect of the “étale-like link” by comparing the “Frobenius-like link” with the *transport* as discussed in (i).



**Corollary 7.7.** — *Let  $(\mathcal{A}, B, \times)$  and  $G \gamma M$  an MLF  $\gamma$ -pair. Then the natural homomorphism*

$$\text{Aut}(G \gamma M) \rightarrow \text{Aut}(G)$$

*fits into the following exact sequence:*

$$1 \rightarrow \text{Ind} \rightarrow \text{Aut}(G \gamma M) \rightarrow \text{Aut}(G) \rightarrow 1.$$

*Proof.* — This assertion is the content of Theorem 7.6 in the case where  $(G \gamma M) = (G \cdot \gamma M \cdot)$ .

### References

- [1] Y. HOSHI, “A note on the geometricity of open homomorphisms between the absolute Galois groups of  $p$ -adic local fields”, *Kodai Math. J.* **36** (2013), no. 2, p. 284-298.
- [2] ———, “Mono-anabelian reconstruction of number fields”, *RIMS Kôkyûroku Bessatsu* **B76** (2019), p. 1-77.
- [3] M. JARDEN & J. RITTER, “On the characterization of local fields by their absolute Galois groups”, *J. Number Theory* **11** (1979), no. 1, p. 1-13.



- [4] S. MOCHIZUKI, "A version of the Grothendieck conjecture for  $p$ -adic local fields", *Int. J. Math.* **8** (1997), no. 4, p. 499-506.
- [5] ———, "Topics in absolute anabelian geometry I: generalities", *J. Math. Sci., Tokyo* **19** (2012), no. 2, p. 139-242.
- [6] ———, "Topics in absolute anabelian geometry III: global reconstruction algorithms", *J. Math. Sci., Tokyo* **22** (2015), no. 4, p. 939-1156.
- [7] ———, "The mathematics of mutually alien copies: from Gaussian integrals to inter-universal Teichmüller theory", *RIMS Kôkyûroku Bessatsu* **B84** (2021), p. 23-192.
- [8] J. NEUKIRCH, *Algebraic number theory*, Grundlehren der Mathematischen Wissenschaften, vol. 322, Springer, 1992.
- [9] J. NEUKIRCH, A. SCHMIDT & K. WINGBERG, *Cohomology of number fields*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer, 2008.
- [10] N. NIKOLOV & D. SEGAL, "Finite index subgroups in profinite groups", *C. R. Math. Acad. Sci. Paris* **337** (2003), no. 5, p. 303-308.
- [11] J.-P. SERRE, "Local class field theory", in *Algebraic Number Theory*, Academic Press Inc., 1965, p. 128-161.
- [12] J. T. TATE, " $p$ -divisible groups", in *Proc. Conf. Local Fields (Driebergen, 1966)*, Springer, 1967, p. 158-183.
- [13] S. YAMAGATA, "A counterexample for the local analogy of a theorem by Iwasawa and Uchida", *Proc. Japan Acad.* **52** (1976), no. 6, p. 276-278.

---

January 31, 2017

YUICHIRO HOSHI, Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, JAPAN  
E-mail : yui chi ro@kuri ms. kyoto-u. ac. j p